

Copyright  
Not for publication  
Quintessence

# Clinical Endodontics

Brooke Blicher, DMD  
Rebekah Lucier Pryles, DMD  
Jarshen Lin, DDS  
Alice Li, DMD





## Dedication

*To John Dresser, whose continued commitment to excellence and the pursuit of knowledge sets the highest bar for what it means to provide exceptional patient-centered care.*



One book, one tree: In support of reforestation worldwide and to address the climate crisis, for every book sold Quintessence Publishing will plant a tree (<https://onetreplanted.org/>).

### Library of Congress Cataloging-in-Publication Data

Names: Blicher, Brooke, author. | Lucier Pryles, Rebekah, author. | Lin, Jarshen, author. | Li, Alice, author.

Title: Clinical endodontics / Brooke Blicher, Rebekah Lucier Pryles, Jarshen Lin, Alice Li.

Description: Batavia, IL : Quintessence Publishing Co. Inc., [2023] | Includes bibliographical references and index. | Summary: "This manual equips clinicians with all the best protocols and knowledge for administering general endodontic care"-- Provided by publisher.

Identifiers: LCCN 2023027796 | ISBN 9781647241759 (spiral bound)

Subjects: MESH: Endodontics--methods | Periodontal Diseases | Dental Pulp Diseases

Classification: LCC RK351 | NLM WU 241 | DDC 617.6/342--dc23/eng/20230726

LC record available at <https://lccn.loc.gov/2023027796>

A CIP record for this book is available from the British Library.  
ISBN: 9781647241759



© 2024 Quintessence Publishing Co, Inc


Quintessence Publishing Co, Inc  
411 N Raddant Road  
Batavia, IL 60510  
[www.quintessence-publishing.com](http://www.quintessence-publishing.com)

5 4 3 2 1

All rights reserved. This book or any part thereof may not be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, or otherwise, without prior written permission of the publisher.

Editor: Kristen Clark  
Design: Sue Zubek  
Production: Angelina Schmelter

Printed in China



# Clinical Endodontics

**Brooke Blicher, DMD**

Private Practice Limited to Endodontics  
White River Junction, Vermont

Lecturer  
Department of Restorative Dentistry  
and Biomaterials Sciences  
Harvard School of Dental Medicine

Assistant Clinical Professor  
Department of Endodontics  
Tufts University School of Dental Medicine  
Boston, Massachusetts

**Jarshen Lin, DDS**

Director of Predoctoral Endodontics  
Department of Restorative Dentistry and  
Biomaterials Sciences  
Harvard School of Dental Medicine

Clinical Associate  
Department of Surgery, Division of Dentistry  
Massachusetts General Hospital  
Boston, Massachusetts

**Rebekah Lucier Pryles, DMD**

Private Practice Limited to Endodontics  
White River Junction, Vermont

Assistant Clinical Professor  
Department of Endodontics  
Tufts University School of Dental Medicine

Lecturer  
Department of Restorative Dentistry  
and Biomaterials Sciences  
Harvard School of Dental Medicine  
Boston, Massachusetts


**Alice Li, DMD**

MMSc Candidate  
Advanced Graduate Program in Endodontics  
Harvard School of Dental Medicine  
Boston, Massachusetts

Chief Content Officer  
My Dental Key, LLC

 **QUINTESSENCE PUBLISHING**

Berlin | Chicago | Tokyo  
Barcelona | London | Milan | Mexico City | Paris | Prague | Seoul | Warsaw  
Beijing | Istanbul | Sao Paulo | Zagreb



# Contents

**Preface** *vi*

**PART I /** Establishing a Diagnosis *1*

**1 Examination Protocols** *3*

**2 The Definitive Diagnosis** *27*

**PART II /** Management of Endodontic  
Pathology *39*

**3 Diagnosis-Driven Treatment  
Planning** *41*

**4 Pharmacology** *49*

**5 Nonsurgical Endodontics** *66*

**6 Nonsurgical Retreatment** *115*

**7 Apical Microsurgery** *125*

**8 Vital Pulp Therapy** *137*



9 **Treatment of Immature Necrotic  
Teeth** 143

10 **Emergency Treatment and Complication  
Management** 152

**PART III /** Adjunctive Endodontic  
Diagnoses and Treatments 177

11 **Fractures** 179

12 **Resorption** 190

13 **Periodontal-Endodontic Infections** 204


14 **Traumatic Dental Injuries** 208

**Glossary** 229

**Index** 237

**QUICK GUIDE**

*Booklet insert located in pocket of back cover.*



# Preface

**E**ndodontic care is integral to the practice of dentistry. Sadly, however, many practitioners are intimidated by endodontic diagnosis and treatment planning. This text aims to cover the full breadth of endodontic diagnosis and care to bring an evidence-based perspective to clinical practice. The vast swath of endodontic literature offers clinicians both classic wisdom and new information that can be applied directly to patient care, and readers will find literature references throughout the text to support evidence-based practice.

Though the origins of this book were in the development of predoctoral endodontics curricula, its comprehensive scope renders it useful for practitioners of all levels, including dental students, residents, general practitioners, and specialists alike. Even providers who do not perform the full scope of endodontic procedures should be knowledgeable about their existence and availability.

The main text is dedicated to the overall theory and biologic basis of diagnosis and treatment, including detailed procedure guides. The Quick Guide, found at the back of the book, is modeled on cookbooks and includes tray setups and step-by-step procedural instructions, making this guide useful for not only practitioners but also clinical staff tasked with operator setup.

We hope you enjoy our illustrated and evidence-based guide to clinical endodontic practice and that it promotes both your learning and the delivery of excellent clinical care.



## PART I

# Establishing a Diagnosis

The practice of endodontics is grounded in the management of orofacial pain and infection. Pain and infection, however, are not exclusive to endodontic pathology. Furthermore, not all endodontic diagnoses warrant endodontic treatment. The establishment of an accurate and complete diagnosis is an essential prerequisite for any endodontic treatment. This section of the text discusses the materials and techniques for detecting and diagnosing endodontic pathology, as well as its differential diagnosis.




# Examination Protocols



**D**iagnosis is the foundation of endodontics. The diagnosis of endodontic pathology and orofacial pain requires gathering a careful patient history and performing both clinical and radiographic examinations. These components are referred to as the subjective and objective exams. Taken together, the exam findings are used to establish the patient's diagnosis and direct the selection and provision of appropriate treatment modalities. This chapter reviews the components of the diagnostic exam.

## Subjective Examination

As is true of all medical and dental encounters, a thoughtful and thorough patient interview, also known as the subjective examination, is both foundational to the development of a trusting patient-clinician relationship and provides necessary information to direct the objective examination that follows. This interview can take place either in person or, when in-person encounters are not possible, as a telehealth encounter via phone or video as permitted by local practice laws. The components of the subjective exam are (1) the chief complaint, (2) the history of the present illness (HPI), (3) the past dental history (PDH), and (4) the medical history (Fig 1-1).



**FIG 1-1** The components of the subjective examination.

The chief complaint is the reason the patient is seeking care. This should be recorded in the patient's own words, not only to demonstrate to the patient that their concerns are understood but also to ensure that treatment plans address the patient's explicit and individualized needs.

The HPI should further develop the clinician's understanding of the patient's chief complaint. If not immediately volunteered, the timeline of symptoms (ie, the onset and duration) and their trajectory over time (increased, decreased, or steady) should be established. Furthermore, clinicians must inquire about symptom intensity, localization, and exacerbating and alleviating factors. If analgesic use is reported, the timing of the last dosage should be established because medications may directly affect responses to clinical testing.

Because endodontic pathology develops as a result of pulpal irritation, the PDH provides clues to the etiology of the chief complaint, including but not limited to restorative care, endodontic treatment, fractures, and trauma. The setting, type, and timing of prior dental care should be ascertained. If trauma is included in the history, the specifics surrounding the traumatic injury must be understood. Deep restorative care may result in the development of endodontic pathology years after treatment.<sup>1</sup> Teeth with a history of deep restorations face a lifetime of elevated risk for requiring nonsurgical root canal therapy (NSRCT), with the risk being further increased for teeth serving as abutments for partial dentures than for those supporting single crowns.<sup>2-6</sup>

As with all dental examinations, a thorough medical history, including past and present illnesses, medications, and allergies, should be obtained. The medical history can alert clinicians to medications and conditions that may interfere with endodontic diagnosis. Ibuprofen taken to treat the chief complaint or another painful condition can directly mitigate responses to clinical testing, including responses to percussion, palpation, and cold testing.<sup>7</sup> Additionally, a history of head/neck radiation may lessen the expected responses to pulp sensitivity testing.<sup>8</sup>

The medical history may also alert clinicians to other medications or conditions that could impact the delivery of care. Blood thinners warrant consideration both because of their direct effects in increasing bleeding in cases of planned surgical care and because they contraindicate the use of nonsteroidal anti-inflammatory drug (NSAID) family pain relievers.<sup>9,10</sup> Antiresorptive medications, including bisphosphonate and RANKL inhibitors, or a history of head/neck radiation therapy affect treatment planning because they increase the risk of osteonecrosis following invasive dental procedures, including surgical endodontic care and extractions.<sup>11-13</sup> Patients with poorly controlled diabetes or other systemic or drug-related immune system compromise may have difficulty healing from infections, and consideration might be made for antibiotic coverage in consultation with their physicians.<sup>14</sup> Cardiac conditions warranting antibiotic premedication for endodontic procedures should be documented.<sup>15,16</sup>

Measurement of vital signs should also be included in the review of a patient's medical history. This should include measurement of the patient's blood pressure and body temperature. An oral or temporal temperature is important for detecting systemic effects of infection. A high-quality thermometer should be used and the measurement recorded in the patient's chart whenever infection is suspected.


The results of the subjective exam should be recorded in the patient's chart. From a medico-legal perspective, this ensures that patient needs and desires are being communicated and addressed. Most importantly, the patient interview should develop the clinician's differential diagnosis. The interview and differential diagnosis should then be utilized to guide the objective examination that follows.

## Objective Examination

The objective examination includes both a clinical and radiographic examination. The clinical exam consists of a careful extraoral and intraoral inspection of the hard and soft tissues, as well as clinical testing. The radiographic examination should include both 2D and 3D radiographs, when relevant. The standard armamentarium for the objective endodontic exam is shown in Fig 1-2, and the clinical components are detailed in Fig 1-3.



**FIG 1-2** The standard armamentarium for the clinical examination includes a clean mirror, an explorer, a periodontal probe, cotton-tipped applicators or pliers with a no. 2 or no. 4 cotton pellet, refrigerant spray, and bite testing implements including cotton rolls and/or commercially available bite testers, such as a Tooth Slooth (Professional Results) or Fracfinder (Denbur).




**FIG 1-3** The components of the objective examination.

### Extraoral examination

The extraoral examination of the head and neck should assess for swelling, asymmetries, and lymphadenopathy (Fig 1-4). The quality of any swelling should be assessed as fluctuant or firm. Palpation of submandibular and cervical lymph nodes can be used to


## Examination Protocols

detect lymphadenopathy resulting from infections of the head and neck region, though not necessarily specific to the dentition. Extraoral sinus tracts, though rare, should also be documented. These represent a means of drainage from an infected tooth. As with intraoral sinus tracts, they should be radiographically traced with gutta-percha to confirm the source.



**FIG 1-4** Extraoral swelling (a) may indicate an endodontically derived cellulitis, such as this case that occurred secondary to pulpal necrosis with acute apical abscess of the maxillary left first molar (b).

The muscles of mastication and the temporomandibular joint should also be inspected during the extraoral examination (Fig 1-5). Palpation of the masseter and temporalis muscles, particularly in patients with a history of bruxism, is essential to evaluate whether a patient's chief complaint is the result of myofascial pain as opposed to endodontic pathology.



**FIG 1-5** The muscles of mastication and the temporomandibular joint should be examined during the extraoral exam to rule out myofascial pain or a temporomandibular joint disorder as the etiology of the chief complaint.

### Adjunctive trauma examination

When evaluating traumatic dental injuries, the objective examination should be expanded if the clinician is the first to examine the patient following the injury. A primary patient survey should be conducted following the mnemonic **ABCDE**. Clinicians must ensure a patent **airway** and stabilized cervical spine. They should also ensure that there is adequate **breathing** and ventilation and intact **circulation** without evidence of shock. Clinicians must assess for neurologic **disability**, and the patient should be fully undressed to achieve **exposure** of the full body for examination.<sup>17</sup> Additionally, because dental trauma carries the risk of concomitant head trauma, a neurologic screening tool, such as the Glasgow Coma Scale,<sup>18,19</sup> should be used to rule out neurologic compromise or decompensation, which would warrant immediate EMS referral (Table 1-1).

**TABLE 1-1** The Glasgow Coma Scale

Behavior	Response	Score
Eye Opening Response	Spontaneously	4
	To speech	3
	To pain	2
	No response	1
Best Verbal Response	Oriented to time, place, and person	5
	Confused	4
	Inappropriate words	3
	Incomprehensible sounds	2
	No response	1
Best Motor Response	Obeys commands	6
	Moves to localized pain	5
	Flexion withdrawal from pain	4
	Abnormal flexion (decorticate)	3
	Abnormal extension (decerebrate)	2
	No response	1
Total Score	Best response	15
	Comatose client	8 or less
	Totally unresponsive	3

## Intraoral examination

### Hard tissue examination

A comprehensive examination of the dentition is crucial to establish both a potential etiology for endodontic pathology and to evaluate the restorability of the tooth or teeth in question. New or recurrent carious lesions should be identified. Assessment of the lesions with a dental explorer not only discloses caries depth but also reveals any associated sensitivity indicative of near or frank exposures of the adjacent pulp. The overall quality of existing restorations should be evaluated because poorly sealed restorations can permit coronal leakage even in the absence of frank caries.

Cracks or fractures should be visualized and explored for loss of tooth structure and separation or mobility between fractured segments. Cracks and fractures may be better visualized with the use of a fiber optic light to transilluminate the fracture line itself; the fracture will stop transmission of the light, delineating a clear break.<sup>20,21</sup> Certain dyes, including vegetable-based versions (eg, To Dye For, Roydent) and methylene blue, can also be used to more clearly delineate coronal fracture lines<sup>21</sup> (Fig 1-6). It can be difficult to assess the depth of unseparated fracture lines running mesiodistally in posterior teeth due to extensions into interproximal spaces. That said, this assessment is crucial to determine the prognosis for a tooth (see chapters 11 and 14 for more on fractured teeth).



**FIG 1-6** Dyes such as methylene blue or vegetable-based dyes can be used to better visualize fracture lines.



**FIG 1-7** (*a and b*) The assessment of restorability should begin during the examination process because restorability dictates whether endodontic treatment can or should be provided in the presence of pulpal and periapical pathology. (Photo courtesy of Dr Alicia Willette.)

Whether caries or fractures are noted, clinicians must carefully evaluate restorability (Fig 1-7). Deep caries may violate biologic width, warranting consideration of adjunctive procedures, including crown-lengthening surgery or orthodontic extrusion. Cracks and fractures may similarly violate periodontal structures, warranting adjunctive procedures or extraction.

Just as caries and fractures must be assessed, so too must the color of the tooth or teeth in question. Discoloration of a tooth as compared to neighboring controls can indicate transient pulpal pathology or necrosis. Other discolorations may be present because of endodontic or restorative materials within the crown of the tooth, and although these may not create biologic issues, esthetic concerns may warrant management (Fig 1-8). The patient's level of concern about such discoloration will impact treatment planning, including the potential need for internal bleaching following endodontic treatment. Gray or brown discolorations suggest staining secondary to an infected pulp, whereas pink discolorations suggest resorptive etiologies.<sup>22</sup>



**FIG 1-8** Discoloration of the dentition should be assessed as part of every endodontic evaluation. Pink discoloration (*a*) may indicate the development of a resorptive defect, whereas yellow, gray, or brown discoloration (*b*) points to pulp necrosis or dental materials as the cause of discoloration.

A basic examination of occlusal patterns and contacts can be performed to check for alternative sources of pain as well as potential risk factors for fractures. Bruxism and parafunctional habits can both cause initial endodontic pathology when fractures develop and may cause posttreatment failures if not corrected. The presence of wear facets can similarly indicate parafunction and potential occlusal trauma.

It should be noted that there is a clear etiology for most cases of endodontic pathology. That said, spontaneous pulp necrosis may rarely occur due to adjacent nonodontogenic masses or tumors or surgical care leading to devitalization of roots. Case reports have additionally suggested that pulp necrosis might occur secondary to medical conditions including herpes zoster<sup>23</sup> and sickle cell anemia.<sup>24</sup>


### Soft tissue examination

The intraoral soft tissues should be examined during every dental examination. The endodontic exam specifically aims to visualize any signs of intraoral swelling, sinus tracts, or periodontal defects.


- Swelling:** Acute endodontic infections are associated with the development of swelling (Fig 1-9). Many intraoral swellings are visible, and manual palpation can also help to detect subtle swellings, especially when making comparisons to contralateral tissues. The location of the swelling must be carefully documented. Swelling associated with endodontic pathology is often detectable buccal or facial to the tooth, but palatal or lingual swellings may also develop. Endodontically derived swelling is most often located adjacent to the root apex of the affected tooth. Swelling located closer to the gingival margin raises suspicion of periodontal pathology, fractures, or resorptive defects but may also result from periodontal-endodontic infections.<sup>25</sup> As with extraoral swellings, the quality of the swelling as fluctuant or firm should be noted, as well as its dimensions and extension.
- Sinus tracts:** Chronic endodontic infections are associated with sinus tracts. These sinus tracts present as an opening through the alveolar mucosa and may develop on the buccal or lingual/palatal aspects of teeth (Fig 1-10). Like swellings, sinus tracts of endodontic origin are typically located adjacent to the root apex of the affected tooth.<sup>26</sup> Coronally positioned sinus tracts raise suspicion of vertical root fractures or periodontal-endodontic infections. Because sinus tracts may not always present directly adjacent to their source, radiographic tracing with gutta-percha is essential. CBCT imaging represents an alternative radiographic means of assessing sinus tracts; the pathway of bone loss can be followed from the radiographic lesion to the sinus tract opening as visualized clinically.



**FIG 1-9** Localized swelling associated with endodontic pathology typically occurs adjacent to the apices of the teeth involved. This patient presented with an acute apical abscess associated with the maxillary left first molar. The fluctuant swelling was noted apical to the tooth. (Case courtesy of Dr Coco Lin.)



**FIG 1-10** Sinus tracts indicate a draining infection, and radiographs must be used to confirm their source. Both periapical imaging utilizing a gutta-percha cone for tracing or CBCT imaging to accurately show the pathology and its tract through bone to soft tissue are acceptable options. In the case shown, a palatal sinus tract (a) was traced to the maxillary right first molar first with gutta-percha and periapical imaging (b) and then via CBCT imaging without tracing (c).



**FIG 1-11** A limited periodontal exam, including an assessment of probing depths and mobility, should be performed on all teeth undergoing endodontic evaluation.


- **Periodontal exam:** The soft tissue exam should include a limited periodontal exam, incorporating measurement of circumferential probing depths, detection of associated bleeding or purulence, assessment of mobility, and checking for recession (Fig 1-11). Localized narrow and deep probing measurements are associated with both periodontal-endodontic infections and vertical root fractures.<sup>26</sup> Wider defects or generalized probing depths, on the other hand, are more frequently associated with periodontal disease. Bleeding or purulence on probing can indicate inflammation or infection, respectively. Mobility of a single tooth may indicate severe periodontal disease or a larger endodontic infection with loss of bone support. Mobility of several teeth together suggests alveolar fracture.

### Clinical testing

The diagnosis of endodontic pathology requires an accurate replication of the patient's chief complaint. This is accomplished through clinical tests referred to as pulp sensitivity and periapical tests. These tests act as conduits to determine the health of the dental pulp and to detect signs of

inflammation of the adjacent periodontal ligament (PDL).<sup>27</sup> The results should be considered as a whole. It is rare that the results of a single test can be used to confer the absolute diagnosis for a tooth. The results of two or more of these tests will confirm the presence or absence of endodontic disease.

As with all types of diagnostic tests, control testing is essential. These results must also be recorded in the patient's chart. Referred pain from endodontic pathology is common, and confirming the source of pain via clinical and radiographic exam findings is essential. Referral patterns include pain originating in neighboring teeth or even teeth in the opposing arch.<sup>28</sup> When the pain is difficult to localize, testing should be completed on all teeth in the suspected arch as well as the opposing arch. Contralateral teeth may be used as controls when neighboring controls are unavailable or perhaps have been previously endodontically treated. Consideration should be given to the type of restoration on control teeth because teeth with indirect restorations or metallic restorations may globally exhibit stronger responses to thermal testing than those with full-coverage restorations, particularly ceramic restorations. Clinicians must never ignore the possibility that more than one tooth may be the source of symptoms. Figure 1-12 provides an example of control selection.



**FIG 1-12** An example of control teeth used when testing for suspected pathology in the mandibular right first molar. At the very least, the adjacent second molar and second premolar should be tested as control teeth. In patients with poorly localized pain, the area of testing should be expanded to include the remaining teeth in the quadrant (teeth 25 to 28), as well as teeth in the opposing arch (teeth 2 to 8) to rule out referred pain. The mandibular left first molar can also be used as a control tooth; pulp sensitivity tests and PDL tests should provoke similar responses in the contralateral tooth.

### PULP SENSITIVITY TESTS

Testing modalities that can be used to ascertain the health or disease of the pulp tissue include cold testing, heat testing, and electric pulp testing. Pulp sensitivity testing measures a response to conduit measures of nerve sensibility. Presently, no assessment methods for the true measures of pulp vitality are available for use in routine clinical



**FIG 1-13** The armamentarium for pulp sensitivity testing should include a refrigerant spray (eg, EdgeIce, EdgeEndo) and either a cotton-tipped applicator or cotton pellet.

practice, but modalities are under investigation in laboratory settings. For now, the term "pulp sensitivity testing" is more accurate than "pulp vitality testing."

As a result of testing limitations and the fact that the presence of vital tissues does not always correspond to a response to tests, false positive and false negative results are possible. Limitations in accuracy must be considered whenever inconsistent test results arise, and a diagnosis should never be made based solely on the results of sensitivity testing. Pulp sensitivity tests are notoriously inaccurate in immature teeth<sup>29</sup> and immediately following traumatic dental injuries.<sup>30</sup> Consequently, extra care must be taken when making a diagnosis with these comorbidities.

Cold testing is the most accurate pulp sensitivity test currently available.<sup>31</sup> Refrigerant sprays are considered the safest and most convenient and effective means of cold testing.<sup>31,32</sup> These can be

delivered to the surface of the tooth using either cotton-tipped applicators or a cotton pellet held with cotton pliers (Fig 1-13). Because the buccal or facial surface balances the needs for accessibility and proximity to the pulp chamber, it is the surface of choice for cold testing. That said, the occlusal and lingual or palatal surfaces can be tested when an initial response is not detected. A common issue with cold testing is insufficient application of the refrigerant spray to the cotton, which may not elicit a response. Thus, it is important to ensure that the delivery device is sufficiently soaked in cold spray, which may require several seconds of spraying to enhance test accuracy.

Patients should be instructed to raise their hand to communicate the sensation of cold or pain resulting from the test and to keep their hand raised until the sensation diminishes to allow for detection of both the presence and duration of the response. Additionally, patients should be asked to compare the intensity of the sensation felt after each tooth is tested in order to determine if a tooth exhibits a heightened or reduced response compared to controls. Although cold testing may be less accurate on teeth restored with full-coverage restorations,<sup>33</sup> testing should still be completed because many heavily restored teeth will continue to exhibit a response.<sup>34</sup> Ultimately, comparison with similarly restored controls allows for the best assessment of what is normal for the individual patient.

Although heat testing is not as accurate as cold testing,<sup>31</sup> it is useful when heat sensitivity is part of the patient's chief complaint. Heat sensitivity is most commonly reported in teeth with symptomatic irreversible pulpitis or necrosis and has also been reported in previously treated teeth due to the presence of untreated anatomy.<sup>35</sup> Heated gutta-percha is the safest and most effective means for heat testing.<sup>36</sup> This can be done by heating

day or up to several weeks later depending on the situation. In certain cases, reimaging may be undertaken, but considering the principles of ALARA, a reasonable interval of time should pass before exposing the patient to additional radiation to confirm likely radiographic changes.


## References

1. Abou-Rass M. The stressed pulp condition: An endodontic-restorative diagnostic concept. *J Prosthet Dent* 1982;48:264–267.
2. Yavorek A, Bhagavatula P, Patel K, Szabo A, Ibrahim M. The incidence of root canal therapy after full coverage restorations: A 10-year retrospective study. *J Endod* 2020;46:605–610.
3. Valderhaug J, Jokstad A, Ambjørnsen E, Norheim PW. Assessment of the periapical and clinical status of crowned teeth over 25 years. *J Dent* 1997;25:97–105.
4. Saunders WP, Saunders EM. Prevalence of periradicular periodontitis associated with crowned teeth in an adult Scottish subpopulation. *Br Dent J* 1998;185:137–140.
5. Kontakiotis EG, Filippatos CG, Stefopoulos S, Tzanetakakis GN. A prospective study of the incidence of asymptomatic pulp necrosis following crown preparation. *Int Endod J* 2015;48:512–517.
6. Cheung GSP, Lai SCN, Ng RPY. Fate of vital pulps beneath a metal-ceramic crown or a bridge retainer. *Int Endod J* 2005;38:521–530.
7. Read JK, McClanahan SB, Khan AA, Lunos S, Bowles WR. Effect of ibuprofen on masking endodontic diagnosis. *J Endod* 2014;40:1058–1062.
8. Gupta N, Grewal MS, Gairola M, Grewal S, Ahlawat P. Dental pulp status of posterior teeth in patients with oral and oropharyngeal cancer treated with radiotherapy: 1-year follow-up. *J Endod* 2018;44:549–554.
9. Lin S, Hoffman R, Nabriski O, Moreinos D, Dummer PMH. Management of patients receiving novel anti-thrombotic treatment in endodontic practice: Review and clinical recommendations. *Int Endod J* 2021;54:1754–1768.
10. Kaplovitch E, Dounaevskaia V. Treatment in the dental practice of the patient receiving anticoagulation therapy. *J Am Dent Assoc* 2019;150:602–608.
11. Katz H. Endodontic implications of bisphosphonate-associated osteonecrosis of the jaws: A report of three cases. *J Endod* 2005;31:831–834.
12. Ruggiero SL, Fantasia J, Carlson E. Bisphosphonate-related osteonecrosis of the jaw: Background and guidelines for diagnosis, staging and management. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2006;102:433–441.
13. Chrcanovic BR, Reher P, Sousa AA, Harris M. Osteoradionecrosis of the jaws—A current overview—Part 1. *Oral Maxillofac Surg* 2010;14:3–16.
14. Fouad AF, Burleson J. The effect of diabetes mellitus on endodontic treatment outcome: Data from an electronic patient record. *J Am Dent Assoc* 2003;134:43–51.
15. Otto CM, Nishimura RA, Bonow RO, et al. 2020 ACC/AHA Guideline for the management of patients with valvular heart disease: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation* 2021;143:e72–e227 [erratum 2021;143:e229].
16. Wilson WR, Gewitz M, Lockhart PB, et al. Prevention of Viridans group streptococcal infective endocarditis: A scientific statement from the American Heart Association. *Circulation* 2021;143:e963–e978.
17. Steelman R. Rapid physical assessment of the injured child. *J Endod* 2013;39(3 Suppl):S9–S12.
18. Teasdale G, Jennett B. Assessment and prognosis of coma after head injury. *Acta Neurochir (Wien)* 1976;34:45–55.
19. Teasdale G, Jennett B. Assessment of coma and impaired consciousness. A practical scale. *Lancet* 1974;2:81–84.
20. Friedman J, Marcus MI. Transillumination of the oral cavity with use of fiber optics. *J Am Dent Assoc* 1970;80:801–809.
21. Wright HM Jr, Loushine RJ, Weller RN, Kimbrough WF, Waller J, Pashley DH. Identification of resected root-end dentinal cracks: A comparative study of transillumination and dyes. *J Endod* 2004;30:712–715.
22. Heithersay GS. Clinical, radiologic, and histopathologic features of invasive cervical resorption. *Quintessence Int* 1999;30:27–37.
23. Rauckhorst AJ, Baumgartner JC. Zebra. XIX. Part 2. Oral herpes zoster. *J Endod* 2000;26:469–471.
24. Costa CPS, Thomaz EBAF, Souza SFC. Association between sickle cell anemia and pulp necrosis. *J Endod* 2013;39:177–181.
25. Berman LH, Rotstein I. Diagnosis. In: Berman LH, Hargreaves KH (eds). *Cohen's Pathways of the Pulp*, ed 12. St. Louis: Elsevier, 2020:1–33.
26. Rivera E, Walton RE. *Cracking the cracked tooth code: Detection and treatment of various longitudinal tooth fractures*. Chicago: American Association of Endodontics, 2008.

27. Ricucci D, Loghin S, Siqueira JF Jr. Correlation between clinical and histologic pulp diagnoses. *J Endod* 2014;40:1932–1939.
28. Bender IB. Pulpal pain diagnosis—A review. *J Endod* 2000;26:175–179.
29. Fulling HJ, Andreasen JO. Influence of maturation status and tooth type of permanent teeth upon electrometric and thermal pulp testing. *Scand J Dent Res* 1976;84:286–290.
30. Bhaskar SN, Rappaport HM. Dental vitality tests and pulp status. *J Am Dent Assoc* 1973;86:409–411.
31. Mainkar A, Kim SG. Diagnostic accuracy of 5 dental pulp tests: A systematic review and meta-analysis. *J Endod* 2018;44:694–702.
32. White J, Cooley RL. A quantitative evaluation of thermal pulp testing. *J Endod* 1977;3:453–457.
33. Hazard ML, Wicker C, Qian F, Williamson AE, Teixeira FB. Accuracy of cold sensibility testing on teeth with full-coverage restorations: A clinical study. *Int Endod J* 2021;54:1008–1015.
34. Miller SO, Johnson JD, Allemang JD, Strother JM. Cold testing through full-coverage restorations. *J Endod* 2004;30:695–700.
35. Keir DM, Walker WA 3rd, Schindler WG, Dazey SE. Thermally induced pulpalgia in endodontically treated teeth. *J Endod* 1991;17:38–40.
36. Bierma MM, McClanahan S, Baisden MK, Bowles WR. Comparison of heat-testing methodology. *J Endod* 2012;38:1106–1109.
37. Wilson BL, Broberg C, Baumgartner JC, Harris C, Kron J. Safety of electronic apex locators and pulp testers in patients with implanted cardiac pacemakers or cardioverter/defibrillators. *J Endod* 2006;32:847–852.
38. Ketterl W. Age-induced changes in the teeth and their attachment apparatus. *Int Dent J* 1983;33:262–271.
39. Seltzer S, Bender IB, Nazimov H. Differential diagnosis of pulp conditions. *Oral Surg Oral Med Oral Pathol* 1965;19:383–391.
40. Owatz CB, Khan AA, Schindler WG, Schwartz SA, Keiser K, Hargreaves KM. The incidence of mechanical allodynia in patients with irreversible pulpitis. *J Endod* 2007;33:552–556.
41. AAE and AAOMR joint position statement: Use of cone beam computed tomography in endodontics 2015 update. *Oral Surg Oral Med Oral Pathol Oral Radiol* 2015;120:508–512.
42. AAE and AAOMR joint position statement: Use of cone beam computed tomography in endodontics 2015 update [editorial]. *J Endod* 2015;41:1393–1396.
43. Brynolf I. Roentgenologic periapical diagnosis. II. One, two or more roentgenograms? *Sven Tandlak Tidsskr* 1970;63:345–350.
44. Robinson D, Goerig AC, Neaverth EJ. Endodontic access: An update, Part I. *Compendium* 1989;10:290–292, 294–296, 298.
45. American Association of Endodontists. Recommended Guidelines of the American Association of Endodontists for the Treatment of Traumatic Dental Injuries. 2013. [https://www.aae.org/specialty/wp-content/uploads/sites/2/2019/02/19\\_TraumaGuidelines.pdf](https://www.aae.org/specialty/wp-content/uploads/sites/2/2019/02/19_TraumaGuidelines.pdf). Accessed 16 March 2023.
46. Bender IB, Seltzer S. Roentgenographic and direct observation of experimental lesions in bone: II. 1961. *J Endod* 2003;29:707–712.
47. Vande Voorde HE, Bjorndahl AM. Estimating endodontic “working length” with paralleling radiographs. *Oral Surg Oral Med Oral Pathol* 1969;27:106–110.
48. Chang L, Umorin M, Augsburger RA, Glickman GN, Jalali P. Periradicular lesions in cancellous bone can be detected radiographically. *J Endod* 2020;46:496–501.
49. Gutmann JL, Endo C. Clark’s Rule vis a vis the buccal object rule: Its evolution & application in endodontics. *J Hist Dent* 2011;59:12–15.
50. Patel S, Dawood A, Mannocci F, Wilson R, Pitt Ford T. Detection of periapical bone defects in human jaws using cone beam computed tomography and intraoral radiography. *Int Endod J* 2009;42:507–515.
51. Lofthag-Hansen S, Huumonen S, Gröndahl K, Gröndahl HG. Limited cone-beam CT and intraoral radiography for the diagnosis of periapical pathology. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2007;103:114–119.
52. Low KMT, Dula K, Bürgin W, von Arx T. Comparison of periapical radiography and limited cone-beam tomography in posterior maxillary teeth referred for apical surgery. *J Endod* 2008;34:557–562.
53. Ludlow JB, Ivanovic M. Comparative dosimetry of dental CBCT devices and 64-slice CT for oral and maxillofacial radiology. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2008;106:106–114.
54. Ludlow JB, Timothy R, Walker C, et al. Effective dose of dental CBCT—A meta analysis of published data and additional data for nine CBCT units. *Dentomaxillofac Radiol* 2015;44:20140197.
55. Pauwels R, Beinsberger J, Collaert B, et al. Effective dose range for dental cone beam computed tomography scanners. *Eur J Radiol* 2012;81:267–271.
56. Farman AG. ALARA still applies. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2005;100:395–397.
57. Patel S, Brown J, Semper M, Abella F, Mannocci F. European Society of Endodontology position statement: Use of cone beam computed tomography in endodontics: European Society of Endodontology (ESE) developed by. *Int Endod J* 2019;52:1675–1678.

58. Fayad MI. The impact of cone beam computed tomography in endodontics: A new era in diagnosis and treatment planning. American Association of Endodontics, 2018. <https://www.aae.org/specialty/news-letter/the-impact-of-cone-beam-computed-tomography-in-endodontics-a-new-era-in-diagnosis-and-treatment-planning>. Accessed 16 March 2023.
59. Costa FF, Gaia BF, Umetsubo OS, Cavalcanti MGP. Detection of horizontal root fracture with small-volume cone-beam computed tomography in the presence and absence of intracanal metallic post. *J Endod* 2011;37:1456–1459.
60. Scarfe WC, Levin MD, Gane D, Farman AG. Use of cone beam computed tomography in endodontics. *Int J Dent* 2009;2009:634567.
61. Setzer FC, Shi KJ, Zhang Z, et al. Artificial intelligence for the computer-aided detection of periapical lesions in cone-beam computed tomographic images. *J Endod* 2020;46:987–993.
62. Lam EWN, Mallya SM. *White and Pharaoh's Oral Radiology: Principles and Interpretation*, ed 8. St Louis: Mosby, 2018.
63. Brynolf I. *A Histological and Roentgenological Study of the Periapical Region of Human Upper Incisors* [thesis]. Stockholm: Almqvist and Wiksell, 1967.
64. Peters E, Lau M. Histopathologic examination to confirm diagnosis of periapical lesions: A review. *J Can Dent Assoc* 2003;69:598–600.
65. Lalonde ER. A new rationale for the management of periapical granulomas and cysts: An evaluation of histopathological and radiographic findings. *J Am Dent Assoc* 1970;80:1056–1059.
66. Weissman J, Johnson JD, Anderson M, et al. Association between the presence of apical periodontitis and clinical symptoms in endodontic patients using cone-beam computed tomography and periapical radiographs. *J Endod* 2015;41:1824–1829.
67. Sener S, Cobankara FK, Akgünlü F. Calcifications of the pulp chamber: Prevalence and implicated factors. *Clin Oral Investig* 2009;13:209–215.
68. Pettiette MT, Wright JT, Trope M. Dentinogenesis imperfecta: Endodontic implications. Case report. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 1998;86:733–737.
69. Pettiette MT, Zhong S, Moretti AJ, Khan AA. Potential correlation between statins and pulp chamber calcification. *J Endod* 2013;39:1119–1123.
70. Gold SI. Root canal calcification associated with prednisone therapy: A case report. *J Am Dent Assoc* 1989;119:523–525.
71. Sayegh FS, Reed AJ. Calcification in the dental pulp. *Oral Surg Oral Med Oral Pathol* 1968;25:873–882.
72. Edds AC, Walden JE, Scheetz JP, Goldsmith LJ, Drisko CL, Eleazer PD. Pilot study of correlation of pulp stones with cardiovascular disease. *J Endod* 2005;31:504–506.
73. Robertson A, Andreasen FM, Bergenholtz G, Andreasen JO, Norén JG. Incidence of pulp necrosis subsequent to pulp canal obliteration from trauma of permanent incisors. *J Endod* 1996;22:557–560.
74. Blicher B, Pryles RL. The use of selective anesthesia in endodontic diagnosis. *Compend Contin Educ Dent* 2021;42:498–575.
75. White JJ, Reader A, Beck M, Meyers WJ. The periodontal ligament injection: A comparison of the efficacy in human maxillary and mandibular teeth. *J Endod* 1988;14:508–514.

# Index



Page numbers followed by “t” denote tables; those followed by “f” denote figures.

## A

AAE. *See* American Association of Endodontists.

### Abscess

- acute periodontal, 32, 32f
- apical. *See* Apical abscess.

### Access

- in nonsurgical retreatment, 116
- in nonsurgical root canal therapy. *See* Nonsurgical root canal therapy, access in.

Acetaminophen, 49–50, 50t

Acute periodontal abscesses, 32, 32f

ALARA principle, 18, 24, 106

### Alveolar bone

- fracture of, 218, 218f, 225
- infection of, 32

Ameloblastoma, 33f

### American Association of Endodontists

- antibiotics prescribing guidelines of, 155
- Case Difficulty Assessment Form, 47, 155
- Glossary of Endodontic Terms, 179, 179f, 181, 212, 213f

Amoxicillin, 52, 53f, 53t, 55t

Amoxicillin with clavulanic acid, 52, 53f, 53t

Ampicillin, 55t

Analgesics, 4, 49–50, 50t

### Anesthesia

- adjunctive techniques for, 60–63
- apical microsurgery, 127–128
- confirmation of, 60, 60f
- intraosseous injections for, 61, 61f
- intraosseous injections for, 62, 62f
- intrapulpal injections for, 62–63, 63f
- local, 57, 60t, 127–128, 128f
- mandibular, 59, 59f, 60t
- maxillary, 59, 59f, 60t
- pulpal, 57, 59, 60f
- selective, 22–23, 23f, 61
- topical, 58, 58f, 134

Anesthetics, 57–58, 58f

Anterior teeth. *See also* specific teeth.

- definitive restorations in, 102f
- dental dam isolation of, 66
- mandibular, 59
- nonsurgical root canal therapy access preparation in, 72, 73f
- treatment planning in, 41

Antibiotic pastes, 92f, 92–93, 149

Antibiotics. *See also* specific antibiotic.

- adjunctive medical management uses of, 155
- contraindications for, 51, 51t
- description of, 50
- dosing of, 52
- flare-ups managed with, 167
- indications for, 51, 51t–52t
- premedication uses of, 54f–55f, 54–55
- systemic, 51–52

Anticoagulants, 126

Antiresorptive medications, 4

Anxiolytics, 56–57, 68

### Apexification

- bioceramic, 144f, 144–146
- calcium hydroxide, 145–147, 146f–147f
- cervical root fracture treated with, 45
- indications for, 45
- pulpectomy in, 143–144
- purpose of, 143
- risks associated with, 143, 144f
- success rates for, 43t

Apexogenesis, 44, 137, 138f. *See also* Vital pulp therapy.

### Apical abscess

#### acute

- antibiotics for, 51
- examination findings in, 28t
- incision and drainage of, 153, 153f

#### chronic

- antibiotics for, 51
- examination findings in, 28t

Apical constriction, 78f, 78–79, 100

Apical external inflammatory root resorption, 193, 194f

### Apical microsurgery

- anesthesia for, 127–128
- armamentarium for, 128f
- CBCT imaging in, 127, 127f, 130
- definition of, 125
- flaps in, 129–130, 129f–130f, 133
- follow-up after, 135, 135f
- grafting in, 132
- hemostasis in, 128, 128f
- incisions in, 129, 129f
- indications for, 125, 126f
- lead excision and biopsy, 131, 131f
- magnification for, 128
- osteotomy in, 130, 130f
- pain control in, 127–128
- in posterior maxilla, 130
- postoperative care for, 133–134, 134f
- preoperative evaluation for, 126–127, 127f
- for previously treated teeth, 44
- resection in, 131f, 131–132
- retrofilling in, 131–132, 132f
- retropreparation in, 131–132, 132f
- sinus exposure during, 134
- success rates for, 43t
- topical anesthetics in, 134
- transient paresthesia associated with, 168

Apical perforations, 160, 160f

Apical periodontitis, 28t, 51

Apical termination, 78–79

Apical zipping, 157, 157f

Apicomarginal bony defects, 205

Articaine, 57, 59

As low as reasonably achievable principle. *See* ALARA principle.

Athletic mouthguards, 211, 211f

Azithromycin, 53, 53t, 55t

**B**

Bacteremia, transient, 54  
 Bacterial endocarditis, 54  
 Benzodiazepines, 56  
 Bioceramic apexification, 144f, 144–146  
 Bisphosphonates, 4, 126  
 Bite blocks, 68  
 Bite sensitivity testing, 14–15, 15f  
 Bitewing radiographs, 16, 16f, 72, 72f, 161  
 Bleaching, internal, 170–172, 170f–172f  
 Blood thinners, 4  
 Broaches, 82f, 83  
 Brown discoloration, 8, 8f  
 Bruxism, 8  
 Buccal Object Rule, 16  
 Bupivacaine, 58  
 Burs, 70, 70f, 77

**C**

C files, 82f, 83  
 Calcification, 21, 21f  
 Calcium hydroxide  
   apexification uses of, 145–147, 146f–147f  
   description of, 45  
   intracanal medicament uses of, 92, 92f–93f, 123, 149, 165  
   localized tissue necrosis caused by, 165  
   pulp capping uses of, 138  
   sealers, 96, 97f  
 Cardiac pain, 35  
 Cardiac transplants, 54, 55f  
 Caries  
   deep, 8  
   examination for, 7–8, 8f  
   intraoral photography of, 21  
   removal of, 139  
 CBCT imaging  
   advantages of, 17, 17f  
   alveolar fractures on, 218f  
   apical microsurgery uses of, 127, 127f, 130  
   canals located using, 76, 77f  
   cemental tears on, 187, 187f  
   crown-root fractures on, 215, 215f  
   follow-up uses of, 18, 19f, 106, 107f, 135, 135f  
   fractures on, 180, 180f, 183, 184f  
   indications for, 18, 19b  
   internal root resorption on, 192f  
   limitations of, 17, 20  
   maxillary sinusitis of endodontic origin on, 30, 31f  
   orthogonal planes, 18, 18f  
   perforations on, 161, 161f  
   periapical cemento-osseous dysplasia on, 37f  
   periapical pathology on, 29, 30f  
   periodontal ligament luxations on, 221f  
   preoperative, in apical microsurgery, 127, 127f  
   pulp chamber position on, 71, 72f  
   radiation dosage from, 17–18  
   root fractures on, 216, 216f  
   sinus tracts on, 9, 10f  
   working length determinations using, 79  
 Cefazolin, 55t  
 Ceftriaxone, 55t  
 CEJ. *See* Cementoenamel junction.  
 Cellulitis, 7f  
 Cemental tears, 187, 187f  
 Cementoblastoma, 37f  
 Cementodentinal junction, 78, 78f  
 Cementoenamel junction, 70, 75, 117, 150, 170  
 Cephalixin, 53, 53t  
 Cephalosporins, 55

Cervical lymph nodes, 5  
 Chemomechanical preparation, 81  
 Chief complaint, 4, 10, 27  
 Chlorhexidine gluconate, 68, 87f, 89, 92, 127  
 Chloroform, 119, 119f  
 Circulation, 6  
 Clark's Rule, 16, 17f  
 Clindamycin, 53, 53f, 53t  
 Clinical testing  
   control testing in, 11, 11f  
   documentation of findings, 15t  
   overview of, 10–11  
   periapical tests, 14–15, 14f–15f  
   pulp sensitivity tests, 11–14, 12f–13f, 29f, 183, 185, 187, 209, 218  
 Coalescing primary endodontic and periodontal lesions, 206–207, 207f  
 Cold lateral compaction, 98, 98f, 100  
 Cold testing, of pulp, 12, 12f, 60  
 Complete pulpotomy, 141  
 Complicated crown fractures, 181, 213t, 214–215, 215f  
 Complications  
   apical zipping, 157, 157f  
   flare-ups, 167, 167f  
   informed consent discussions regarding, 155  
   instrument separation, 157–159, 158f–159f  
   internal bleaching for discoloration, 170–172, 170f–172f  
   ledges, 156, 156f  
   material extrusion beyond the apex, 164–165, 168  
   overview of, 155–156  
   paresthesia, 168, 168f  
   perforations, 159–163, 160f–162f  
   persistent pain, 168–170, 169f  
   sodium hypochlorite accident, 163f–164f, 163–164  
   thermal injury, 165–167, 166f  
   transportation, 157, 157f  
 Concussion, of periodontal ligament, 219t, 220, 220f  
 Condensing osteitis, 20, 20f, 28t  
 Cone fit radiographs, 97, 97f  
 Continuous wave of condensation, 99  
 Coronal disassembly, 117, 117f  
 Coronal flaring, 85  
 Coronal fractures, 21  
 Corticosteroids, 169  
 Cracked tooth syndrome, 185  
 Cracks, 7–8, 8f, 179  
 Craze lines, 181  
 Crestal perforations, 162  
 Crown fractures  
   complicated, 213t, 214–215, 215f  
   description of, 179, 181, 210  
   uncomplicated, 213t, 213–215  
 Crown-down technique, 86–87, 87f  
 Crown-root fractures, 179, 181, 215, 215f

**D**

Decoronation, 194, 195f, 225  
 Deeper sedation, 57  
 Definitive restorations, 100, 103, 103f  
 Dental dam isolation  
   armamentarium for, 67f  
   for direct pulp capping, 139  
   for indirect pulp capping, 139  
   for nonsurgical root canal therapy, 66–68, 67f  
   for pulpotomy, 141  
   for regenerative endodontics, 150  
 Dentoalveolar pain disorder, 169  
 Diabetes mellitus, 4, 126  
 Diagnosis  
   adjunctive, 30  
   of endodontic pathology, 27–31

- inability to replicate symptoms for, 23–24  
of nonendodontic pathology, 32–37  
objective examination for. *See* Objective examination.  
periapical, 27, 27f  
pulpal, 27, 27f  
requirements for, 3  
selective anesthesia for, 22–23, 23f  
subjective examination for, 3f, 3–5  
testing on another day, 23–24  
treatment planning based on, 41–47
- Direct osteotomy, 130
- Direct pulp capping, 139–140, 140f
- Disability, 6
- Discolorations  
coronal, 214, 215f  
description of, 8, 8f  
internal bleaching for, 170–172, 170f–172f
- Double-ended endodontic explorer, 77, 77f
- Doxycycline, 55t
- Drains, 153
- Dycal, 138, 140
- Dyes, 7–8, 8f, 131
- E**
- EALs. *See* Electronic apex locators.
- ECR. *See* External cervical resorption.
- EDTA, 87f, 88–90, 144, 149f, 150
- EIRR. *See* External inflammatory root resorption.
- Electric pulp testing, 13f, 13–14
- Electronic apex locators, 78–80, 79f, 85, 156, 161, 217
- Emergency treatment  
adjunctive medical management, 155, 155f  
description of, 152  
incision and drainage, 153, 153f  
infection management in, 155f  
modalities for, 153f  
pain management in, 155  
pulpectomy, 154–155  
pulpotomy, 154, 154f  
triage, 152f
- EndoActivator, 91, 91f
- Endocarditis, 54
- Endodontics  
complications of. *See* Complications.  
goal of, 78  
success rates for, 43t
- Epinephrine, 58, 127–128
- EPT. *See* Electric pulp testing.
- Examination  
extraoral, 5–6  
objective. *See* Objective examination.  
subjective, 3f, 3–5
- External cervical resorption, 191, 196–200, 201t, 196f–200f
- External inflammatory root resorption, 193, 193f–194f, 195, 201t
- Extraoral examination, 5–6
- Extraoral sinus tracts, 6
- Extrusive luxation, of periodontal ligament, 219t, 220f, 220–221
- F**
- Files  
fracture of, 158f  
hand, 81f–82f, 81–83, 119  
rotary, 83f–85f, 83–85, 119, 157
- Finger spreaders, 98, 98f
- Flaps, 129–130, 129f–130f, 133
- Flare-ups, 167, 167f
- Follow-up  
apical microsurgery, 135, 135f  
CBCT imaging for, 18, 19f, 106, 107f, 135, 135f  
nonsurgical root canal therapy, 106, 106f–107f
- Formocresol, 140
- Fractures  
alveolar, 218, 218f, 225  
CBCT imaging of, 183, 184f  
crown. *See* Crown fractures.  
crown-root, 179, 181, 213t, 215f, 215–216  
dyes for visualizing, 7–8, 8f, 180  
examination for, 7–8, 8f  
root. *See* Root fractures.  
traumatic, 179, 181, 212–218, 213t, 214f–217f  
2D imaging of, 180, 180f  
unseparated, 184f–185f, 184–186  
vertical root, 10, 179, 182, 183f, 184
- Full pulpotomy, 43t, 154
- G**
- Gates Glidden instruments, 84f, 84–85, 119
- Glasgow Coma Scale, 6, 7t
- Glide path, 85, 159
- Gutta-percha  
in bioceramic apexification, 145  
heat testing uses of, 12–13  
obturation uses of, 94–96, 95f, 98, 100f  
removal of, 103, 119–120, 170
- H**
- Hand files, 81f–82f, 81–83, 119
- Hard tissue examination, 7–9, 8f
- Head and neck examination, 5–6
- Headaches, 35
- Heat testing, of pulp, 12–13, 13f, 29
- Hedstrom files, 82f, 83, 119–120, 120f
- Heithersay's classification system, 198f
- Hemostasis  
in apical microsurgery, 128, 128f  
in incision and drainage, 153
- History of present illness, 3–4
- Horizontal root fractures, 181f, 181–182, 217f
- I**
- Iatrogenic perforations, 159–161
- Ibuprofen, 23, 49–50, 50t
- Idiopathic peri-cementitis, 169
- Immature necrotic teeth  
apexification for. *See* Apexification.  
description of, 45, 143  
regenerative endodontics for, 147–150, 147f–151f
- Immunocompromised patients, 51
- Incision and drainage, 153, 153f
- Indirect pulp capping, 138–139, 139f
- Infections  
alveolar bone, 32  
nonodontogenic, 32, 33f  
periodontal, 32, 32f  
periodontal-endodontic, 204–207, 204f–207f
- Inferior alveolar nerve block, 56–58
- Inflammation  
anesthesia in, 58  
periapical tests for, 14
- Informed consent, 45–47, 46f, 104, 155, 167
- Infractions, 181

Instrumentation  
 nonsurgical root canal therapy. *See* Nonsurgical root canal therapy, instrumentation.  
 separation of, 157–159, 158f–159f  
 ultrasonic, 158, 158f–159f  
 Internal bleaching, 170–172, 170f–172f  
 Internal root resorption, 191–192, 191f–192f, 201t, 212f  
 Intra canal medicaments  
 description of, 91–93, 92f–93f  
 extrusion of, beyond the apex, 164–165, 168  
 Intraligamentary injections, 61, 61f  
 Intraoral examination  
 clinical testing in  
 control testing in, 11, 11f  
 documentation of findings, 15t  
 overview of, 10–11  
 periapical tests, 14–15, 14f–15f  
 pulp sensitivity tests, 11–14, 12f–13f, 29f, 183, 185, 187, 209, 218  
 hard tissue, 7–9, 8f  
 periodontal examination, 10, 10f  
 soft tissue, 9f, 9–10  
 Intraoral photography, 21, 22f  
 Intraorifice barriers, 101, 101f, 186  
 Intraosseous injections, 62, 62f  
 Intrapulpal injections, 62–63, 63f  
 Intrasulcular incision, 129, 129f  
 Intrusive luxation, of periodontal ligament, 219t, 222f, 222–223, 223t  
 IRM retrofillings, 125, 125f  
 IRR. *See* Internal root resorption.  
 Irreversible pulpitis  
 antibiotic contraindications in, 51  
 asymptomatic, 28t, 43, 137, 141f  
 examination findings in, 28t  
 nonsurgical root canal therapy for, 44, 137  
 pulpectomy for, 154  
 symptomatic, 28t, 186, 206  
 vital pulp therapy for, 43–44  
 Irrigation  
 in access preparation, 74  
 active, 90–91, 91f  
 armamentarium, 87–89  
 in bioceramic apexification, 144  
 chlorhexidine gluconate, 87f, 89  
 delivery devices, 89, 89f  
 EDTA, 87f, 88–90, 144, 149f, 150  
 methods of, 90–91  
 protocol for, 90f  
 in regenerative endodontics, 149f  
 side-vented needles used in, 89, 89f  
 sodium hypochlorite, 87f–88f, 88–91, 150

**J**

Jaw radiolucencies, 36, 36f

**K**

K files, 82, 82f  
 Krasner and Rankow's laws  
 for canal location, 76, 76f  
 for pulp chamber location, 70, 71f

**L**

Lateral condensation, 98f, 98–99  
 Lateral external inflammatory root resorption, 193, 194f  
 Lateral luxation, of periodontal ligament, 219t, 220f, 221–222

Ledges, 156, 156f  
 Lesions of endodontic origin, 20, 20f  
 Local anesthesia  
 description of, 57, 60t, 127–128, 128f  
 paresthesia caused by, 168  
 Longitudinal root fractures, 182–183  
 Loop instruments, 158, 159f  
 Loupes  
 for apical microsurgery, 128  
 for nonsurgical root canal therapy, 68–69, 69f  
 Luxation injury, 209  
 Lymph nodes, 5–6  
 Lymphadenopathy, 6

**M**

MAF. *See* Master apical file.  
 Magnification  
 for apical microsurgery, 128  
 for instrument removal, 158  
 for nonsurgical root canal therapy, 68–69, 69f  
 for perforation prevention, 163  
 Maltingering, 36  
 Mandibular anesthesia, 59, 59f, 60t  
 Mandibular block anesthesia, 22  
 Mandibular incisors, 75t  
 Mandibular molars  
 anesthesia of, 59  
 canals/roots in, 75t  
 ledges on, 156f  
 nonsurgical root canal therapy access preparation in, 74  
 Mandibular premolars  
 anesthesia of, 59  
 canals/roots in, 75t  
 description of, 73  
 necrotic, 150f  
 Masseter muscle, 6, 6f  
 Master apical file, 80, 86, 86f  
 Mastication muscles, 6, 6f  
 Maxillary anesthesia  
 description of, 59, 59f, 60t  
 selective anesthesia, 22, 23f  
 Maxillary canines, 75t  
 Maxillary incisors, 75t  
 Maxillary molars  
 acute apical abscess of, 7f  
 canals/roots in, 75t  
 nonsurgical root canal therapy access preparation in, 74  
 Maxillary premolars  
 canals/roots in, 75t  
 description of, 73  
 perforation in, 161f  
 Maxillary sinusitis of endodontic origin, 30, 31f  
 Medical history, 3–4, 56, 126, 127f  
 Medication-related osteonecrosis of the jaw, 33f, 126  
 Medications, 4. *See also* Pharmacology; specific medication.  
 Mepivacaine, 61  
 Methylene blue, 7, 8f, 131  
 Metronidazole, 52–53, 53f, 53t  
 Micromirrors, 131f, 131–132  
 Microsurgery. *See* Apical microsurgery.  
 Migraine headaches, 35  
 Mineral trioxide aggregate  
 apical barrier creation using, 100  
 as retrofilling, 132  
 staining caused by, 170  
 Modified Masserann technique, 121, 122f, 158  
 Molars  
 mandibular. *See* Mandibular molars.  
 maxillary. *See* Maxillary molars.  
 nonsurgical root canal therapy access preparation in, 74, 74f

- MRONJ. See Medication-related osteonecrosis of the jaw.
- MSEO. See Maxillary sinusitis of endodontic origin.
- MTA. See Mineral trioxide aggregate.
- Multivisit therapy
- for nonsurgical retreatment, 122–123
  - for nonsurgical root canal therapy, 91–92, 92f
  - for regenerative endodontics, 148
- Myofascial pain, 34
- N**
- Nasopalatine duct cyst, 36f
- Neuropathic pain, 35
- Neurovascular pain, 35
- Nickel-titanium rotary instruments
- description of, 81–84, 85f, 87
  - separation of, 157–158
- Nitrous oxide, 56–57, 68
- Nonodontogenic infections, 32, 33f
- Nonodontogenic pain, 34f, 34–36
- Nonodontogenic swelling, 32
- Nonsteroidal anti-inflammatory drugs, 49–50, 50t, 155
- Nonsurgical endodontics, 66
- Nonsurgical retreatment
- access in, 116
  - candidates for, 116
  - coronal disassembly in, 117, 117f
  - definition of, 115
  - indications for, 44, 116, 116f
  - multivisit, 122–123
  - obturation material removal in, 119–122, 119f–123f
  - pathology requiring, 115–116, 116f
  - post removal in, 117–118, 118f
  - for previously treated teeth, 44
  - prior treatment corrected by, 122
  - single-visit, 122–123
  - success rates for, 43t, 44
- Nonsurgical root canal therapy
- access in
    - anterior teeth, 72, 73f
    - armamentarium for, 69f–70f, 69–72
    - bitewing radiographs of, 72, 72f
    - bleeding during, 74
    - burs for, 70, 70f
    - canal location for, 77
    - challenges associated with, 74–75
    - depth of, 72, 72f
    - intracoronar restoration removal during, 70
    - irrigation during, 74
    - molars, 74, 74f
    - premolars, 73, 74f
    - pulp chamber influence on, 70, 71f
    - purpose of, 69
    - radiographic anatomy for, 71, 72f
    - shape of, 70–72, 71f
    - straight-line, 71–72, 72f–73f
    - temporary crown removal for, 70
  - apical termination measurements, 78–79
  - apicomarginal bony defects treated with, 205
  - armamentarium, 66, 67f, 69f–70f, 69–72, 81–85, 81f–85f
  - canal location in
    - access dependence on, 77
    - CBCT imaging of, 76, 77f
    - double-ended endodontic explorer for scouting of, 77, 77f
    - tactile sensation used for, 77–78
  - canal morphology, 75t
  - chemomechanical preparation, 81
  - dental dam isolation for, 66–68, 67f
  - discomfort after, 104–105
  - electronic apex locators in, 78–80, 79f, 85
  - flare-ups after, 105, 105f
  - follow-up appointments, 106, 106f–107f
  - indications for, 41, 41f
  - informed consent for, 46, 46f, 104
  - instrumentation
    - armamentarium for, 81, 81f
    - crown-down technique, 86–87, 87f
    - hand files, 81f–82f, 81–83, 119
    - methods of use, 85–87, 86f–87f
    - rotary files, 83f–85f, 83–85, 119
    - step-back technique, 86, 86f
  - internal root resorption treated with, 192, 192f
  - intracanal medicaments used in, 91–93, 92f–93f
  - irreversible pulpitis treated with, 44, 137
  - irrigation
    - in access preparation, 74
    - active, 90–91, 91f
    - armamentarium, 87–89
    - chlorhexidine gluconate, 87f, 89
    - delivery devices, 89, 89f
    - EDTA, 87f, 88–90
    - methods of, 90–91
    - protocol for, 90f
    - side-vented needles used in, 89, 89f
    - sodium hypochlorite, 87f–88f, 88–91
  - loupes for, 68–69, 69f
  - magnification for, 68–69, 69f
  - multivisit, 91–92, 92f
  - obturation in
    - apical barrier, 100
    - armamentarium, 94–97, 95f, 97f
    - cold lateral compaction, 98, 98f
    - cone fit radiographs for, 97, 97f
    - description of, 91, 93–94
    - drying of canals before, 94, 94f
    - gutta-percha for, 94–96, 95f, 98
    - material for, 93–94, 94f
    - methods, 97–100
    - sealers for, 95–97, 97f
    - 2D imaging of, 191f
    - warm vertical condensation, 99–100, 99f–100f, 119
  - postoperative care for, 104–106, 104f–106f
  - posttreatment pathology, 115
  - preoperative rinses in, 68
  - pulp chamber in, 70, 71f, 76
  - pulp necrosis treated with, 44, 186
  - restorative care
    - definitive, 100, 103, 103f
    - overview of, 100–101
    - posts, 102–103, 103f
    - temporary, 101, 101f
    - timing of, 104
  - risk factors for, 4
  - single-visit, 91, 92f
  - success with, 42f, 43t
  - traumatic fractures treated with, 181
  - working length determination for, 78f–79f, 78–80
- Nonvital bleaching, 21
- NSAIDs. See Nonsteroidal anti-inflammatory drugs.
- NSRCT. See Nonsurgical root canal therapy.
- O**
- Objective examination
- adjunctive trauma examination in, 6, 7f
  - armamentarium for, 5, 5f
  - components of, 5f
  - extraoral examination, 5–6
  - intraoral examination. See Intraoral examination.
  - intraoral photography, 21, 22f
  - radiographic examination. See Radiographic examination.

Obturation  
 apical barrier, 100  
 armamentarium, 94–97, 95f, 97f  
 cold lateral compaction, 98, 98f  
 cone fit radiographs for, 97, 97f  
 description of, 91, 93–94  
 drying of canals before, 94, 94f  
 materials for  
   carrier-based, 94, 120–121  
   chemical removal of, 119, 119f  
   extrusion of, beyond the apex, 164–165, 168  
   gutta-percha, 94–96, 95f, 98  
   mechanical removal of, 119, 119f  
   paresthesia caused by, 168  
   pastes, 94, 121, 121f  
   removal of, 119–122, 119f–122f  
   silver point, 121–122  
 methods, 97–100  
 sealers for, 95–97, 97f  
 thermal injury caused by, 165–166  
 2D imaging of, 191f  
 warm vertical condensation, 99–100, 99f–100f, 119

Occlusal contacts, 8  
 Occlusal pattern, 8  
 Occlusal reduction, 78  
 Odontogenic infections, 32, 32f  
 Odontogenic pain, 33–34, 34f  
 Opioids, 50  
 Oral temperature, 4  
 Osteonecrosis  
   description of, 4  
   medication-related osteonecrosis of the jaw, 33f  
 Osteoperiostitis, periapical, 30, 31f  
 Osteotomy, 130, 130f

**P**

Pain  
 in apical microsurgery, 127–128  
 cardiac, 35  
 emergency treatment management of, 155  
 myofascial, 34  
 neuropathic, 35  
 neurovascular, 35  
 nonodontogenic, 34f, 34–36, 169  
 odontogenic, 33–34, 34f, 169  
 persistent, 168–170, 169f  
 persistent idiopathic facial, 35  
 psychogenic, 36  
 referred, 11  
 vascular, 35

Panoramic radiographs, 16, 16f  
 Papilla-preserving incision, 129, 129f  
 Parafunctional habits, 8  
 Paresthesia, 168, 168f  
 Partial pulpotomy, 43t, 141, 154  
 Past dental history, 3–4  
 Pathologic perforations, 159–160  
 Patient interview, 3–4  
 PCO. *See* Pulp canal obliteration.  
 PDAP. *See* Persistent dentoalveolar pain disorder.  
 PDL. *See* Periodontal ligament.  
 Penicillin VK, 52–53, 53f, 53t  
 Percussion testing, 14, 14f  
 Perforations, 159–163, 160f–162f  
 Periapical cemento-osseous dysplasia, 37f  
 Periapical diagnoses  
   CBCT imaging of, 29, 30f  
   list of, 27, 27f

  pulpal diagnoses and, 29f  
   subjective and objective findings, 28t

Periapical mucositis, 31f  
 Periapical osteoperiostitis, 30, 31f  
 Periapical radiographs, 15  
 Periapical tests, 14–15, 14f–15f  
 Pericanalar resorption resistant sheet, 197, 197f  
 Periodontal disease, 181f  
 Periodontal examination, 10, 10f, 180  
 Periodontal infections, 32, 32f  
 Periodontal ligament  
   concussion of, 219t, 220, 220f  
   electronic apex locators with, 79  
   extrusive luxation of, 219t, 220f, 220–221  
   injuries to, 218–223, 219t, 220f–222f, 223t  
   intraalveolar injection of, 61, 61f  
   intrusive luxation of, 219t, 222f, 222–223  
   lateral luxation of, 219t, 220f, 221–222  
   subluxation of, 219t, 220, 220f  
   widening of, 20, 219t, 221f

Periodontal-endodontic infections, 204–207, 204f–207f  
 Periradicular bony defects, 183  
 Persistent dentoalveolar pain disorder, 35  
 Persistent idiopathic facial pain, 35  
 Persistent pain, 168–170, 169f

Pharmacology  
   analgesics, 49–50, 50t  
   anesthetics, 57–58  
   antibiotics. *See* Antibiotics.  
   anxiolytics, 56–57, 68

Photography, intraoral, 21, 22f  
 PIFP. *See* Persistent idiopathic facial pain.

Post(s)  
   fiber, 102, 118, 118f  
   longitudinal fractures and, 182f  
   metal, 118f  
   nonthreaded, 103  
   removal of, 117–118, 118f  
   selection of, 102–103, 103f

Posterior teeth. *See also* specific teeth.  
   cuspal-coverage restorations in, 102  
   treatment planning in, 41–42

PR. *See* Pressure resorption.

Precementum, 190, 193  
 Preentin, 190  
 Premedication, 54f–55f, 54–55

Premolars  
   dental dam isolation of, 66  
   mandibular. *See* Mandibular premolars.  
   maxillary. *See* Maxillary premolars.  
   nonsurgical root canal therapy access preparation in, 73, 74f

Preoperative rinses, 68  
 Pressure resorption, 195f, 195–196, 201t

Previously initiated teeth  
   endodontic treatments for, 44  
   subjective and objective findings in, 44

Previously treated teeth  
   nonsurgical retreatment for, 44  
   subjective and objective findings in, 44

Probing depths, 10, 10f  
 Psychogenic pain, 36

Pulp  
   intraalveolar injection effects on, 61  
   periodontium and, 204f  
   regeneration of, 147–150, 147f–151f

Pulp canal obliteration, 21, 21f  
 Pulp capping  
   direct, 139–140, 140f

indirect, 138–139, 139f  
 success rates for, 43t

Pulp chamber  
 coronal disassembly, 117  
 in nonsurgical root canal therapy, 70, 71f, 76

Pulp necrosis  
 antibiotic contraindications in, 51, 52t  
 chief complaint in, 27, 28t  
 crown fractures with, 214  
 dental trauma as cause of, 145f  
 discoloration as sign of, 8  
 examination findings in, 28t  
 illustration of, 7f  
 in immature teeth. *See* Immature necrotic teeth.  
 nonsurgical root canal therapy for, 44, 186  
 spontaneous, 9

Pulp sensitivity tests, 11–14, 12f–13f, 29f, 183, 185, 187, 209, 218

Pulpal anesthesia, 57, 59, 60f

Pulpal diagnoses  
 list of, 27, 27f  
 periapical diagnoses and, 29f  
 subjective and objective findings, 28t

Pulpal irritation, 4

Pulpectomy, 143–144, 154–155

Pulpitis  
 bleeding associated with, 161  
 examination findings in, 28t  
 irreversible. *See* Irreversible pulpitis.  
 reversible. *See* Reversible pulpitis.

Pulpotomy, 43t, 140–141, 141f, 154, 154f

## R

Radiographic examination  
 bitewing radiographs, 16, 16f, 161  
 CBCT imaging. *See* CBCT imaging.  
 description of, 15  
 interpretation, 20–21  
 lesions of endodontic origin on, 20, 20f  
 nonendodontic pathology, 36–37, 36f–37f  
 panoramic radiographs, 16, 16f  
 perforations diagnosed with, 161  
 periapical radiographs, 15  
 radiopacities, 36, 37f  
 3D imaging, 17f–19f, 17–20, 161  
 2D imaging, 15–16, 16f–17f, 135, 161  
 working length determinations, 79f, 79–80

Radiopacifiers, 146

Radiopacities, 36, 37f

Radix entomolaris, 76, 77f

RANKL inhibitors, 4

Reamers, 82f, 83

Rectangular flap, 129, 129f

Referred pain, 11

Refrigerant sprays, 12, 12f

Regenerative endodontics, 147–150, 147f–151f

Regenerative therapy, 45

Replacement resorption, 193–195, 194f, 201t, 209, 226, 226f

Replantation, 224–226, 227t

Resin-based sealers, 96, 97f

Resin-modified glass-ionomer temporary restorations, 101, 101f

Resorbable membranes, 132

Resorption  
 description of, 190, 190f  
 external cervical, 191, 196–200, 201t, 196f–200f  
 external inflammatory root, 193, 193f–194f, 195, 201t  
 internal root, 191–192, 191f–192f, 201t  
 pressure, 195f, 195–196, 201t  
 replacement, 193–195, 194f, 201t, 209, 226, 226f

Restorative care  
 definitive, 100, 103, 103f  
 overview of, 100–101  
 posts, 102–103, 103f  
 temporary, 101, 101f

Retractors, 130, 130f

Retrofills, 131–132, 132f

Reversible pulpitis  
 causes of, 42  
 description of, 206  
 examination findings in, 28t  
 illustration of, 186f  
 treatment planning for, 42

Rinses, preoperative, 68

Root fractures  
 CBCT imaging of, 216, 216f  
 description of, 179, 181f–183f, 181–184  
 horizontal, 181f, 181–182, 217f  
 longitudinal, 182–183  
 vertical, 10, 179, 182, 183f, 184

Root perforations, 165

Root resorption  
 external inflammatory, 193, 193f–194f, 195, 201t  
 internal, 191–192, 191f–192f, 201t, 212f

Rotary files, 83f–85f, 83–85, 119, 157

RR. *See* Replacement resorption.

## S

Same Lingual Opposite Buccal Rule, 16, 17f

SAMPLE mnemonic, 208, 209f

Sealer puffs, 165, 165f

Sealers, 95–97, 97f, 165

Sedation, deeper, 57

Selective anesthesia, 22–23, 23f, 61

Side-vented needles, 89, 89f, 164, 164f

Silver point obturation materials, 121–122

Simple interrupted sutures, 133, 133f

Single tooth isolation, 67, 68f

Single-visit therapy  
 for nonsurgical retreatment, 122–123  
 for nonsurgical root canal therapy, 91, 92f

Sinus tracts  
 CBCT imaging of, 9, 10f  
 coronally positioned, 9  
 extraoral, 6  
 intraoral photography of, 22f  
 presentation of, 9, 10f  
 vertical root fractures associated with, 183, 183f

Sinusitis  
 maxillary sinusitis of endodontic origin, 30, 31f  
 nonodontogenic, 34

Sling sutures, 133

SLOB Rule. *See* Same Lingual Opposite Buccal Rule.

Sodium hypochlorite  
 accidents involving, 163f–164f, 163–164  
 irrigation uses of, 87f–88f, 88–91, 95, 139, 149f, 150, 162

Sodium perborate, 171, 172f

Soft tissue  
 examination of, 9f, 9–10  
 lacerations of, 226  
 necrosis of, 166  
 palpation of, 14  
 swelling of, 9, 9f

Split fractures, 182


Spontaneous pupal necrosis, 9

Spreaders, 98, 98f, 100

Staining, of teeth, 170–172, 170f–172f

Stem cells, 147

Step-back technique, 86, 86f



Strip perforations, 160, 160f, 162  
 Subgingival perforations, 162  
 Subjective examination, 3f, 3–5  
 Subluxation, of periodontal ligament, 219t, 220, 220f  
 Submandibular lymph nodes, 5  
 Submarginal incision, 129, 129f  
 Superoxol, 171  
 Supracrestal perforations, 162  
 Surface resorption, 193  
 Surgical operating microscope, 69, 78, 118, 158  
 Swelling  
     examination of, 9, 9f  
     nonodontogenic, 32  
     percussion testing for, 14, 14f  
     sodium hypochlorite accident as cause of, 163, 163f  
 Symptoms, inability to replicate, 23–24

**T**

Teeth  
     discoloration of, 8, 8f  
     fractures of. *See* Fractures.  
     percussion testing of, 14, 14f  
     replantation of, 224–226, 227t  
     restorability assessments, 41–42  
     staining of, 170–172, 170f–172f  
 Temporal arteritis, 35  
 Temporalis muscle, 6, 6f  
 Temporary restorations, 101, 101f  
 Temporomandibular joint, 6, 6f  
 Tension-type headaches, 35  
 Thermal injury, 165–167, 166f, 214  
 Thermoplastic injection, 100  
 3D imaging, 17f–19f, 17–20, 161  
 Topical anesthesia, 58, 58f, 134  
 Transient bacteremia, 54  
 Transportation, 157, 157f  
 Traumatic dental injuries  
     adjunctive examination for, 6, 7t  
     alveolar fractures, 218, 218f  
     antibiotic indications in, 51  
     avulsion, 223f–224f, 223–226  
     fractures. *See* Fractures.  
     periodontal ligament, 218–223, 219t, 220f–222f, 223t  
     primary assessment after, 208, 208f  
     pulp necrosis secondary to, 145f  
     treatment prioritization for, 210f  
     workup after, 209–210  
 Traumatic fractures, 179, 181  
 Treatment planning  
     decision tree used in, 42f–43f  
     diagnosis-driven, 41–47  
     informed consent in, 45–47, 46f  
     pulpal diagnosis in, 42  
     restorability of tooth as focus of, 41  
     root maturation stage in, 42

Triangular flap, 129f  
 Tricalcium silicate cement sealers, 96–97  
 Trigeminal autonomic cephalalgias, 35  
 Trigeminal neuralgia, 35  
 Triple antibiotic pastes, 93, 170  
 Troughing, 118  
 Tugback, 93  
 2D imaging, 15–16, 16f–17f, 135, 161, 180, 180f

**U**

Ultrasonic instruments  
     separation of, 158, 158f–159f  
     thermal injury caused by, 165–166  
 Uncomplicated crown fractures, 181, 213t, 213–214, 214f  
 Unseparated fractures, 184f–185f, 184–186

**V**

Vascular pain, 35  
 Vegetable-based dyes, 7, 8f, 131  
 Vertical condensation, 99–100, 99f–100f  
 Vertical root fractures, 10, 179, 182, 183f, 184  
 Vital pulp therapy  
     armamentarium for, 137f  
     bioceramic materials for, 43–44  
     complicated crown fracture treated with, 215f  
     direct pulp capping, 139–140, 140f  
     indications for, 137, 138f  
     indirect pulp capping, 138–139, 139f  
     irreversible pulpitis treated with, 43–44  
     materials used in, 137–138  
     pulpotomy, 140–141, 141f  
     success of, 45  
 Vital pulp tissue, 29  
 Vital signs, 4

**W**

Walking bleach, 170  
 Warm vertical condensation, 99–100, 99f–100f, 119  
 Winged clamps, 67  
 Working length determination  
     ledges and, 156, 156f  
     in nonsurgical root canal therapy, 78f–79f, 78–80  
     in regenerative endodontics, 148, 149f

**Z**

Zinc-oxide eugenol sealers, 96, 97f  
 Zipping, apical, 157, 157f