

Lena Katharina Müller, Julia Heider, Roland Frankenberger, Christian Graetz, Lutz Jatzwauk, Jens Nagaba, Romy Brodt, Rugzan Jameel Hussein, Anke Weber, Kai Voss, Markus Tröltzsch, Bilal Al-Nawas

Guideline: Dealing with aerosol-borne pathogens in dental practices

Introduction: It is well known that droplets and aerosols may cause infections in dental staff [21]. Therefore adequate protective measures against pathogens transmitted via droplets or aerosols from the patients' oral cavity are of great importance in dental practices. Due to close contact between dental professionals and patients' oral cavity and the formation of droplets, spray mist and aerosols during dental interventions, hygiene and precautionary measures are used in dental practice to prevent the transmission of infectious diseases.

Methods: Relevant information regarding the SARS-CoV-2 and COVID-19 pandemic was obtained from electronic databases such as PubMed, Cochrane library, Web of Science, using the following search terms: "SARS-CoV-2" OR "COVID-19", "airborne transmission", "mouth rinse", "dental", "aerosol" OR "aerosol generating procedures", "droplet", "FFP2" OR "FFP3" OR "N95" OR "mask". Latest reports and guidelines from major health authorities such as the Robert Koch-Institut (RKI), Centers for Disease Control and Prevention (CDC), World Health Organization (WHO), as well as major national dental associations and health regulatory bodies were also referred.

Results: Protecting dental professionals and patients from infections while ensuring basic dental care for the population is of paramount importance. With that in mind, this guideline presents recommendations for dental practitioners during the COVID-19 pandemic.

Keywords: droplets; aerosols; infections; COVID-19 pandemic; prevention; dental practice

Department of Oral and Maxillofacial Surgery, Plastic Surgery, University Medical Centre, Johannes Gutenberg-University, Mainz, Germany: Dr. Lena Katharina Müller, MD, DMD; PD Dr. Dr. Julia Heider, MD, DMD; Prof. Dr. Dr. Bilal Al-Nawas, MD, DMD

Department for Operative Dentistry, Endodontics, and Pediatric Dentistry, Philipps University of Marburg and University Hospital Giessen and Marburg Campus Marburg, Germany: Prof. Dr. Roland Frankenberger, DMD

Department of Conservative Dentistry and Periodontology, University of Kiel, Kiel, Germany: PD Dr. Christian Graetz, DMD

Department of Hospital Infection Control, University Hospital, Dresden, Germany: Prof. Dr. Lutz Jatzwauk

German Dental Association (Bundeszahnärztekammer, BZÄK), Berlin, Germany: Dr. Jens Nagaba, DMD

National Association of Statutory Health Insurance Dentists (Kassenzahnärztliche Bundesvereinigung, KZBV), Cologne Germany: Dr. Romy Brodt, DMD; Dr. Rugzan Jameel Hussein, DMD

German Society of Dentistry and Oral Medicine (Deutsche Gesellschaft für Zahn-, Mund- und Kieferheilkunde e.V., DGZMK): Dr. Anke Weber, DMD

private practice, Kirchbarkau and German Working Committee for Hygiene in Dentistry (Deutscher Arbeitskreis für Hygiene in der Zahnmedizin, DAHZ): Dr. Kai Voss, DMD private practice, Ansbach, Germany and German Society of Dentistry and Oral Medicine (Deutsche Gesellschaft für Zahn-, Mund- und Kieferheilkunde e.V., DGZMK): Dr. Dr. Markus Tröltzsch, DMD, MD

Citation: Müller LK, Heider J, Frankenberger R, Graetz C, Jatzwauk L, Nagaba J, Brodt R, Hussein RJ, Weber A, Voss K, Tröltzsch M, Al-Nawas B: Guideline: Dealing with aerosol-borne pathogens in dental practices. Dtsch Zahnärztl Z Int 2020; 2: 240–245

DOI.org/10.3238/dzz-int.2020.0240-0245

Providing basic dental care and ensuring personal protection in dental practices

The World Health Organization (WHO) associates aerosol-generating medical procedures with increased risk of infection for medical staff from SARS-CoV-2 [53]. Depending on the current situation of the pandemic, it is recommended to avoid these procedures if possible. However, aerosols must not be equated with the spray mist that occurs in dentistry. It is generally known that spray mist can contain pathogens, but in a form that is strongly diluted with cooling water. The term aerosol basically defines a suspension of liquid and solid particles with a diameter of 5 µm, deposits and living or dead microorganisms in a gaseous medium [48, 49]. Spray is a droplet mixture of air, water, solids with particles and is visible to the naked eye. A rebound effect of spray occurs after the impact on the tooth or soft tissue, emerges like a bell from the oral cavity in the work area and, in addition to the spray mist, contains germs, abrasive particles, saliva and possibly blood [11, 14]. The transition from “droplets” to “aerosols” and vice versa is smooth and depends on the ambient conditions. Both aerosol and spray can contain transmissible pathogens [3, 23]. The word aerosol is often colloquially used for all of these potentially infectious media for the sake of simplicity. However, it can be assumed that aerosol-generating dental procedures are certainly less infectious than saliva or bronchial secretions due to the high proportion of cooling water. The present guideline explicitly refers only to the formation of spray and aerosols during dental work.

Even if the regional prevalence of SARS-CoV-2 is high, all dental treatments that alleviate the patients' symptoms or prevent an existing disease from worsening must be guaranteed. It is important to differentiate between healthy or asymptomatic patients and suspected or confirmed COVID-19 infected patients, who should only be treated in compliance with special protective measures.

Triage of suspected cases

Suspected cases should be screened by phone or via a notice on the door at

latest prior to the start of any dental treatment, preferably before the patient even enters the practice. Typical symptoms of an infection with SARS-CoV-2 and questions regarding potential contacts with COVID-19 positive patients in the past 2 weeks should specifically be asked. The body temperature might be measured as part of the triage of suspected cases. However, a large number of false positive results must be assumed. In addition, false negative results may occur if SARS-CoV-2 infected people show no signs of fever or antipyretic agents have been used [43].

Entering the dental practice

When entering the practice, patients should be asked to wear a mask covering both mouth and nose until the start of treatment as well as afterwards. Consistent implementation of basic hygiene including hand hygiene is expected. When entering the practice, patients should be asked to wash or disinfect their hands. Depending on the epidemiological situation, magazines, toys and other expendable items might be dispensed within the waiting room [34, 38]. Since transmission via contact surfaces cannot be ruled out, in addition to basic hygiene, regular disinfection of contact surfaces should be carried out [34, 52].

In order to protect risk groups from infection with SARS-CoV-2, the dental treatment should be integrated into the daily routine in a way that there is as little contact as possible with other patients. Suspected and confirmed COVID-19 cases should preferably be treated in special centers, clinics or practices. If this is not possible in exceptional cases, necessary treatments should be carried out in the dental practice in strategic and or scheduled separation from the patients attending regular consultation, while ensuring all hygiene and safety measures specified for this purpose.

Distancing

Patients should be kept at a distance from staff by observing the minimum distance of 1.5 m for registration [34, 38]. Installing plexiglass shields at registration to further protect employees from droplets. The distance between patients from dif-

ferent households should be at least 1.5 m in order to minimize the risk of the infection being transmitted via droplets [34, 38]. Employees should wear surgical masks permanently, even outside treatment rooms, and maintain the minimum distance requirement, also during breaks and in changing rooms [2, 6, 50].

COVID-Testing

Personnel showing symptoms of a COVID-19 infection should be isolated immediately and tested for the presence of an infection using PCR. There is not enough reliable data to routinely test symptom-free employees in dental practices, but it might be useful in case of a heightened risk situation.

Patients who show symptoms of a COVID-19 infection should only be treated in case of emergency until a negative test can be produced. In the event of a dental emergency, emergency treatment should be carried out in compliance with special protective measures.

Dental emergencies in symptomatic and infected patients

If possible, all dental treatments for symptomatic patients or confirmed COVID-19 patients should be postponed to a later date. In case of a dental emergency treatment (pain, abscesses, infections, complications e.g. secondary bleeding, trauma, etc.), the measures as described in table 1. should be applied:

- strict spatial separation from all other patients,
- patients should wear a surgical mask until the start of treatment,
- where possible, schedule emergency treatment at the end of the day,
- maximum PPE
 - (1) safety glasses/face shield
 - (2) FFP2/FFP3 or N95 mask
 - (3) hygienic hand disinfection
 - (4) disposable gloves
 - (5) headgear and socks (to reduce self-contamination)
 - (6) long-sleeved liquid-repellent protective scrubs
- Final cleaning and disinfection of all surfaces with at least limited virucidal surface disinfectants.

| Diameter of the droplet | 0.3 µm | 0.5 µm | 1.0 µm | 5.0 µm | 10 µm |
|-------------------------|-----------------------|-----------------------|----------------------|----------------------|-----------------------|
| Volume of the droplet | 0.014 µm ³ | 0.065 µm ³ | 0.52 µm ³ | 65.5 µm ³ | 523.6 µm ³ |

Table 1 Relationship between volume and diameter of droplets

Aerosol formation in dental practices, protection through surgical masks and treatment cautions

Emission from persons

Droplets are mainly produced by humans when they speak (sing), cough and sneeze. Droplets that are created when speaking, coughing or sneezing range between 1 and > 10 µm in size [54]. The emission of particles containing bacteria acts 400 : 7 : 1 when sneezing : coughing : talking [15, 32, 41]. Droplets larger than 8 µm in size sediment on surfaces immediately, and no later than following a maximum of 20 minutes. With a size of around 4 µm, droplets sediment within 90 minutes. Smaller droplets (aerosols) can remain in the air for up to 30 hours and can then be transmitted over greater distances by air currents [15]. Depending on the relative humidity, droplets can turn into aerosols [7]. When droplets float in the air, they lose water and become so-called droplet nuclei, which are the size of aerosols. In stagnant room air, the size of the droplets reduces from 12–21 µm to around 4 µm within about 10 minutes [51].

The dehydration of droplets can (depending on the respective microorganism) kill or inactivate bacteria and viruses contained in the droplet. Hence the transition from droplets to droplet cores (or the drying out of aerosols) does not necessarily result in further infectivity of the microorganisms contained. Depending on surrounding conditions, the statements of experimental studies on the detection of SARS-CoV-2 viruses in aerosol that are capable of reproducing differ. Virus particles have been found in aerosols in some studies [29, 52]. Whether and how quickly the droplets and aerosols sink or remain suspended in the air depends on the size of the particles as well as a number of

other factors, including temperature and humidity [26]. From the studies up to date, no statement can be made regarding the infectiousness of the virus particles.

Emission from water-cooled dental instruments

With the introduction of high-speed dental preparation instruments, the need for effective cooling of work areas arose in order to avoid thermal damage to the pulp-dentin system. The required amount of liquid for this lies at approx. 50 ml per minute. The liquid is swirled around and partially reflected on various intraoral structures and the instrument itself. Spray mist rebound contains both large liquid droplets and aerosols. The majority of the spray mist rebound consists of droplets ≥ 10 µm [5]. Around 90 % of the larger particles in the dental spray mist with a size of approx. 20 µm fall on the patient's face or body surface [38]. When using a dental turbine at a distance of 10 cm from the oral cavity of the treated patient, the number of particles with a diameter between 0.3 µm and 0.5 µm increased by a factor of 100 and for particles with a diameter of 7 µm by a factor 3 [27]. The number of particles ≥ 10 µm only increased by a factor of 1.7 when the turbine was used at a distance of 20 cm above the oral cavity, as they sediment quickly. Aerosols and droplets that arise during dental treatments are described in the literature with particle sizes of 0.5–20 µm [35, 40]. Due to their low sedimentation speed, aerosols can float several meters away and also infect people in other rooms or people who are in the treatment room at a later point in time [18]. However, the number of virus copies present in liquids, droplets or aerosols is not to be equated with infectious viruses. The exact infection dose required in virus copies

to trigger an infection with SARS-CoV-2 is currently unknown.

Droplets contain significantly more liquid and therefore more microorganisms than aerosols, hence the necessary infectious dose is reached much faster through ingestion of a droplet. The following calculation of the amount of liquid transported in particles of the corresponding size is clear.

Effectiveness of surgical masks and simple textile mouth and nose covers that protect against large particles, as well as “physical distancing” of 1.5 to 2 m as part of the COVID-19 preventive measures indicate that SARS-CoV-2 is mainly transmitted by droplet infections [9, 55]. Both measures only reduce droplets, but not aerosols. Transmission of SARS-CoV-2 by aerosols has also been observed but requires longer contact times with the aerosol (choir samples) with low air exchange and/or increased humidity (slaughtering businesses) in the room in order to achieve the necessary pathogen dose. In dentistry, occurrences of such “super spreading events” are completely absent.

In conclusion, the current evidence base is insufficient to confirm or exclude airborne transmission with SARS-CoV-2 in the context of dental treatments [8, 36]. As such, procedures for reducing the spray mist, consisting of droplets and small, floating particles, represent basic occupational safety measures for the dental team. Since even trained, ergonomically designed dental technology cannot completely prevent the emission of droplets and aerosols from the patients' oral cavity, putting in place additional measures to minimize the transmission of infection becomes inevitable.

Protective effect of face masks

The recommendations of the Commission for Hospital Hygiene and Infection Prevention (KRINKO) at the Ro-

| Type of mask | Minimum retention capacity of the filter with regard to NaCl test aerosol [respectively <i>Staphylococcus aureus</i>] | Maximum permissible total leakage on subjects |
|------------------------------------|--|---|
| FFP 1 | 80 % | 22 % [a] |
| FFP 2 | 94 % | 8 % [a] |
| FFP 3 | 99 % | 2 % [a] |
| NIOSH N 95 | 95 % | 10 % [b] |
| NIOSH N 99 | 99 % | 10 % [b] |
| NIOSH N 100 | 99.97 % | 10 % [b] |
| Medical masks (<i>S. aureus</i>) | [95 %] | Not specified |

Table 2 Comparison of the requirements for particle-filtering half masks and mouth-nose protection (MNS) [13]; [a] Specified for FFP masks with NaCl aerosol in accordance with DIN EN 149 [12]; [b] For NIOSH-N masks derived from the Assigned Protection Factor (APF) of 10 specified by NIOSH. This requires a passed qualitative or quantitative Occupational Safety and Health Administration (OSHA) fit test [19].

(Table 1 and 2: L. Jatzwauk)

bert Koch-Institute are considered state of the art in the prevention of infectious diseases in Germany. In case of respiratory infections or pneumonia caused by coronaviruses (SARS, MERS), the use of an FFP2 mask is recommended. For patients infected with seasonal influenza A or B, one MNS is sufficient. On the other hand, KRINKO recommends a respirator to prevent avian influenza. Patients with open pulmonary tuberculosis should be treated using an FFP2 mask. Patients with open pulmonary tuberculosis caused by multi-resistant *Mycobacterium tuberculosis* (multi-resistant tuberculosis, MDR-Tbc, or extensively resistant tuberculosis, XDR-Tbc) require the wearing of an FFP3 mask with the same pathogen and transmission path. This shows that the recommendations based on a risk analysis are not only influenced by the quality of the “face masks”, but also the clinical consequences to be expected in the event of an infection. The physical and technical testing of respiratory masks is carried out in accordance with DIN EN 149 under practical conditions. Subjects are exposed to an NaCl test aerosol wearing a respirator. The median mass-related particle size of the aerosol is 0.6 micrometers. However, even within these test conditions there is no absolute protection against the inhalation of aerosols (table 2).

Whether this protective effect is also necessary for infectious diseases that are transmitted by much larger droplets from the respiratory tract or by dehydrated aerosols cannot be derived from these model studies.

Recommended use of masks and face shields

The additional use of face protection shields might further increase safety. Dental staff should wear FFP2/FFP3 or N95 masks if contact with patients with suspected or confirmed SARS-CoV-2 infection takes place. During treatment of patients, who are not suspected to be infected with SARS-CoV-2, dental staff should wear a surgical mask. The best possible barrier function is guaranteed through correct fit of the surgical mask (good adjustment in the nose area and maximum lateral tightness). There is currently no reliable data available for the general wearing of an FFP2/FFP3 or N95 mask for all dental activities using water-cooled instruments.

Reusing masks

In the event of supply shortages in connection with COVID-19, mouth-nose protection and FFP/N95 masks might be reused or reprocessed for specific persons. A reasonable approach to reusing masks might be to provide each employee with at least

5 masks and to use them alternately every day, since a possible SARS-CoV-2 contamination of the 4 unused masks is inactivated after 5 days at the latest (European Centre for Disease Prevention and Control). Alternatively, preparation of masks specific to individuals might be carried out. Reprocessing should take place in sterilizers (e.g. at 121 °C), as the method has proven to be effective and gentle on the material [10].

Treatment precautionary measures

Rinsing the mouth or gargling with mucosal antiseptics shortly before dental treatment could briefly reduce potential virus concentrations in the throat and mouth and thus in the spray and aerosol [24]. Clinical studies regarding the reduction of SARS-CoV-2 currently do not exist for any of the mouth rinses listed below. There are indications of limited virucidal effects (against enveloped viruses) for the following antiseptics:

- ≤ 0,1 % Octenidin®
- 1–1,5 % H₂O₂ [38]
- 0.2 % Povidone-Iod [16, 28, 33, 34]
- 0.2 % Chlorhexidin [4, 33, 37]
- 0.2 % Cetylpyridinium Chloride [31]
- ≤ 0.25 % Natriumhypochlorit [20]
- Dequonal® [33]
- Listerine cool mint® [33]

Just before procedures, patients should be asked to rinse their mouth for 30–60 seconds. Further measures to reduce potential virus contamination by droplets and aerosols should be applied in the context of the respective pandemic situation and are listed below. Spray mist extraction system on the treatment unit, used with an effective systematic extraction technique, reduces the spray mist rebound and aerosols by 2/3 [42]. During dental treatments of suspected and confirmed cases, it is recommended to apply all protective measures as listed below.

There are currently no adequate scientific studies on the effectiveness of room air extraction in combination with HEPA filters or disinfection systems to reduce the viral load in dental treatment rooms.

Precautionary measures

If possible, a rubber dam should be installed [1, 11, 34, 38, 47]. Consistent and high-volume evacuation should be guaranteed. Attention should also be paid to a diameter-optimized suction cannula (≥ 10 mm). If this is guaranteed, there is currently no reliable evidence with regards to effectiveness of any additional suction devices [1, 11, 22, 25, 30, 46]. Large-volume spray mist suction should also be used for treatment methods that are carried out without assistance, such as professional tooth cleaning. After treatments in which aerosols have formed, ventilation should be effective [34]. Almost all instruments rotating rapidly or vibrating at high or highest frequency in the dental practice require a cooling medium. Powder-water blasting devices also require a combination of air, liquids and powder to generate the cleaning jet, which is why all these instruments are inherent in the system with a pronounced spray mist formation [1, 34]. Therefore, their use should be avoided in COVID-19 suspected cases, if clinically possible.

Conflicts of interest

Markus Tröltzsch is the author of a book in Quintessence – “Medicine in the dental practice”, published since

23.10.2020, it contains an article on the topic “Corona”. The author Christian Graetz gives credit for having conducted research projects on spray mist/aerosol formation by means of industrial cooperation (study support by Dürr Dental SE, Bietigheim-Bissingen, D; Loser & Co, Leverkusen, D). The other authors declare that there is no conflict of interest as defined by the guidelines of the International Committee of Medical Journal Editors.

References

1. Ather A, Patel B, Ruparel NB, Diogenes A, Hargreaves KM: Coronavirus disease 19 (COVID-19): implications for clinical dental care. *J Endod* 2020; 46: 584–595
2. Bartoszko JJ, Farooqi MAM, Alhazzani W, Loeb M: Medical masks vs N95 respirators for preventing COVID-19 in health-care workers: a systematic review and meta-analysis of randomized trials. *Influenza Other Respir Viruses* 2020; 14: 365–373
3. Bentley CD, Burkhart NW, Crawford JJ: Evaluating spatter and aerosol contamination during dental procedures. *J Am Dent Assoc* 1994; 125: 579–584
4. Bernstein D, Schiff G, Echler G, Prince A, Feller M, Briner W: In vitro virucidal effectiveness of a 0.12 %-chlorhexidine gluconate mouthrinse. *J Dent Res* 1990; 69: 874–876
5. Böhme W, Goldmann L, Regensburger K, Reitemeier B: Untersuchungen zur Ausbreitung des Sprayrückpralls bei unterschiedlichen Arbeitsbedingungen. *Zahn-Mund-Kieferheilkd* 1990; 78: 621–627
6. Chan JF, Yuan S, Zhang AJ et al.: Surgical mask partition reduces the risk of non-contact transmission in a golden Syrian hamster model for Coronavirus Disease 2019 (COVID-19). *Clin Infect Dis* 2020. doi:10.1093/cid/ciaa644
7. Chen LD: Effects of ambient temperature and humidity on droplet lifetime – a perspective of exhalation sneeze droplets with COVID-19 virus transmission. *Int J Hyg Environ Health* 2020 aug; 229: 113568. doi:10.1016/j.ijheh.2020.113568
8. Cheng VCC, Wong SC, Chen JHK et al.: Escalating infection control response to the rapidly evolving epidemiology of the coronavirus disease 2019 (COVID-19) due to SARS-CoV-2 in Hong Kong. *Infect Control Hosp Epidemiol* 2020 Mar 5: 1–6. doi:10.1017/ice.2020.58
9. Chu DK, Akl EA, Duda S, Solo K, Yaacoub S, Schünemann HJ: Physical distancing, face masks, and eye protection to prevent person-to-person transmission of SARS-CoV-2 and COVID-19: a systematic review and meta-analysis. *Lancet* 2020; 395: 1973–1987
10. Darmstadt TU: „Mehrfach eingesetzte Schutzmasken für Klinikpersonal sind sicher“. www.mpa-ifw.tu-darmstadt.de/startseite_mpaifw/aktuelles_detail_mpaifw_387072.de.jsp (last access on 01.10.2020)
11. Day CJ, Sandy JR, Ireland AJ: Aerosols and splatter in dentistry – a neglected menace? *Dent Update* 2006; 33: 601–602, 604–606
12. DIN EN 149: Atemschutzgeräte – Filternde Halbmasken zum Schutz gegen Partikeln – Anforderungen, P. f., Kennzeichnung. Beuth Verlag, Berlin 2001
13. Dreller S, Jatzwauk L, Nassauer A, Paszkiewicz P, Tobys HU, Rüden H: Zur Frage des geeigneten Atemschutzes vor luftübertragenen Infektionserregern. *Gefahrstoffe – Reinhaltung der Luft* 2006; 66: 14–24
14. Drisko CL, Cochran DL, Blieden T et al.: Position paper: sonic and ultrasonic scalers in periodontics. Research, Science and Therapy Committee of the American Academy of Periodontology. *J Periodontol* 200; 71: 1792–1801
15. Duguid JP: The size and the duration of air-carriage of respiratory droplets and droplet-nuclei. *J Hyg* 1946; 44: 471–479
16. Eggers M, Koburger-Janssen T, Eickmann M, Zorn J: In vitro bactericidal and virucidal efficacy of povidone-iodine gargle/mouthwash against respiratory and oral tract pathogens. *Infect Dis Ther* 2018; 7: 249–259
17. European Centre for Disease Prevention and Control. Options for the decontamination and reuse of respirators in the context of the COVID-19 pandemic – 8 June 2020. Stockholm: ECDC; 2020. www.ecdc.europa.eu/en/publications-data/options-decontamination-and-reuse-respirators-covid-19-pandemic (last access on 01.10.2020)
18. Fernstrom A, Goldblatt M: Aerobiology and its role in the transmission of infectious diseases. *J Pathog* 2013: 493960
19. Fit Testing Procedures (Mandatory) – 1910.134 App A. Hrsg.: U. S. Department of Labor, Occupational Safety & Health Administration. April 1998/August 2004. www.osha.gov/pls/oshaweb/owadisp.show_document?p_table=STANDARDS&p_id=9780
20. Galván M, Gonzalez S, Cohen CL et al: Periodontal effects of 0.25 % sodium hypochlorite twice-weekly oral rinse. A pilot study. *J Periodontal Res* 2014; 49: 696–702
21. Goldman HS, Hartman KS: Infectious diseases. Their disease, our unease: infectious diseases and dental practice. *Va Dent J* 1986; 63: 10–19

22. Graetz C, Bielfeldt J, Tillner A, Plauermann A, Dörfer CE: Spatter contamination in dental practices – how can it be prevented? *Rev Med Chir Soc Med Nat Iasi* 2014; 118: 1122–1134
23. Harrel SK: Airborne spread of disease – the implications for dentistry. *J Calif Dent Assoc* 2004; 32: 901–906
24. Herrera D, Serrano J, Roldán S, Sanz M: Is the oral cavity relevant in SARS-CoV-2 pandemic? *Clin Oral Invest* 2020; 24: 2925–2930
25. Jacks ME: A laboratory comparison of evacuation devices on aerosol reduction. *J Dent Hyg* 2002; 76: 202–206
26. Ji Y, Qian H, Ye J, Zheng X: The impact of ambient humidity on the evaporation and dispersion of exhaled breathing droplets: A numerical investigation. *Journal of Aerosol Science* 2018; 115: 164–172
27. Jurischka L: Experimentelle Untersuchung zur Eignung von Mehrweggesichtsmasken für die zahnärztliche Tätigkeit sowie Untersuchungen des während zahnärztlicher Behandlungen entstehenden Aerosols. *Dissertationschrift, Technische Universität Dresden*; 2014
28. Kariwa H, Fujii N, Takashima I: Inactivation of SARS coronavirus by means of povidone-iodine, physical conditions and chemical reagents. *Dermatology* 2006; 212 (Suppl. 1): 119–123
29. Lednický JA, Lauzardo M, Fan ZH et al.: Viable SARS-CoV-2 in the air of a hospital room with COVID-19 patients. *medRxiv* 2020. doi:10.1101/2020.08.03.20167395
30. Li RW, Leung KW, Sun FC, Samaranyake LP: Severe acute respiratory syndrome (SARS) and the GDP. Part II: implications for GDPs. *Br Dent J* 2004; 197: 130–134
31. Marui VC, Souto MLS, Rovai ES, Romito GA, Chambrone L, Pannuti CM: Efficacy of preprocedural mouthrinses in the reduction of microorganisms in aerosol: A systematic review. *J Am Dent Assoc* 2019; 150: 1015–1026.e1. doi:10.1016/j.adaj.2019.06.024
32. McCluskey F: Does wearing a face mask reduce bacterial wound infection? A literature review. *Br J Theatre Nurs* 1996; 6: 18–20, 29
33. Meister TL, Brüggemann Y, Todt D et al.: Virucidal efficacy of different oral rinses against severe acute respiratory syndrome coronavirus 2. *J Infect Dis* 2020. doi:10.1093/infdis/jiaa471
34. Meng L, Hua F, Bian Z: Coronavirus disease 2019 (COVID-19): emerging and future challenges for dental and oral medicine. *J Dent Res* 2020. doi:10.1177/0022034520914246
35. Micik RE, Miller RL, Mazzarella MA, Ryge G: Studies on dental aerobiology. I. Bacterial aerosols generated during dental procedures. *J Dent Res* 1969; 48: 49–56
36. Ong SWX, Tan YK, Chia PY et al.: Air, surface environmental, and personal protective equipment contamination by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) from a symptomatic patient. *Jama* 2020; 323: 1610–1612. doi:10.1001/jama.2020.3227
37. Park JB, Park NH: Effect of chlorhexidine on the in vitro and in vivo herpes simplex virus infection. *Oral Surgery, Oral Medicine, Oral Pathology* 1989; 67: 149–153
38. Peng X, Xu X, Li Y, Cheng L, Zhou X, Ren B: Transmission routes of 2019-nCoV and controls in dental practice. *Int J Oral Sci* 2020; 12: 9. doi:10.1038/s41368-020-0075-9
39. Perić R, Perić M: Analytical and numerical investigation of the airflow in face masks used for protection against COVID-19 virus – implications for mask design and usage. doi:org/10.15480/882.2775
40. Prospero E, Savini S, Annino I: Microbial aerosol contamination of dental health-care workers' faces and other surfaces in dental practice. *Infect Control Hosp Epidemiol* 2003; 24: 139–141
41. Ransjö U: Masks: a ward investigation and review of the literature. *Journal of Hospital Infection* 1986; 7: 289–294
42. Reitemeier, B: Effektive Reduktion des Spraynebel-Rückpralls – Möglichkeiten und Grenzen. *ZMK* 2010; 26: 662–673
43. Robert Koch-Institut (RKI, 14.05.2020). Fachliche Einschätzung zur Durchführung von Temperaturmessungen und anderen Methoden im Rahmen von Entry- und Exit-Screening an Flughäfen während der COVID-19-Lage, Deutschland. www.rki.de/DE/Content/Infekt/EpidBull/Archiv/2020/20/Art_02.html (last access on 01.10.2020)
44. RKI: Hinweise zur Testung von Patienten auf Infektion mit dem neuartigen Coronavirus SARS-CoV-2. www.rki.de/DE/Content/InfAZ/N/Neuartiges_Coronavirus/Vorl_Testung_nCoV.html (last access on 11.08.2020)
45. RKI: Steckbrief Coronavirus. www.rki.de/DE/Content/InfAZ/N/Neuartiges_Coronavirus/Steckbrief.html (last access on 01.10.2020)
46. Samaranyake LP, Peiris M: Severe acute respiratory syndrome and dentistry: a retrospective view. *J Am Dent Assoc* 2004; 135: 1292–1302
47. Samaranyake LP, Reid J, Evans D: The efficacy of rubber dam isolation in reducing atmospheric bacterial contamination. *ASDC J Dent Child* 1989; 56: 442–444
48. Schulze-Röbbbecke R, Reska M, Lemmen S: Welche Schutzmaske schützt vor COVID-19? Was ist evidenzbasiert? *Krankenhaushygiene* up2date 2020; 15: 123–132
49. Shiu EYC, Leung NHL, Cowling BJ: Controversy around airborne versus droplet transmission of respiratory viruses: implication for infection prevention. *Curr Opin Infect Dis* 2019; 32: 372–379
50. Smith JD, MacDougall CC, Johnstone J, Copes RA, Schwartz B, Garber GE: Effectiveness of N95 respirators versus surgical masks in protecting health care workers from acute respiratory infection: a systematic review and meta-analysis. *Cmaj* 2016; 188: 567–574
51. Stadnytskyi V, Bax CE, Bax A, Anfinrud P: The airborne lifetime of small speech droplets and their potential importance in SARS-CoV-2 transmission. *Proc Natl Acad Sci U S A* 2020; 117, 11875–11877
52. van Doremalen N, Bushmaker T, Morris DH et al.: Aerosol and surface stability of SARS-CoV-2 as compared with SARS-CoV-1. *N Engl J Med* 2020. doi:10.1056/NEJMc2004973
53. WHO: Transmission of SARS-CoV-2: implications for infection prevention precautions. www.who.int/news-room/commentaries/detail/transmission-of-sars-cov-2-implications-for-infection-prevention-precautions (last access on 01.10.2020)
54. Yang S, Lee GW, Chen CM, Wu CC, Yu KP: The size and concentration of droplets generated by coughing in human subjects. *J Aerosol Med* 2007; 20: 484–494
55. Zhang R, Li Y, Zhang AL, Wang Y, Molina MJ: Identifying airborne transmission as the dominant route for the spread of COVID-19. *Proc Natl Acad Sci U S A* 2020; 117: 14857–14863



(Photo: University Medicine Mainz)

DR. LENA KATHARINA MÜLLER
MD, DMD; Department of Oral and Maxillofacial Surgery, Plastic Surgery, University Medical Centre, Johannes Gutenberg-University, Mainz, Augustusplatz 2, 55131 Mainz, Germany
lena_katharina.mueller@unimedizin-mainz.de